

On-Farm Corn and Soybean Planter Demonstration Trials

RFR-A1745

Jim Fawcett, extension field
agronomist (retired)
Andrew Weaver, Northwest Farm,
ag specialist
Lyle Rossiter, Allee Farm, superintendent
Chris Beedle, Western Farm, superintendent
Matthew Schnabel, Northern Farm,
superintendent
Zack Koopman, Ag Engineering/Agronomy
Farm, ag specialist
Cody Schneider, Southeast Farm, ag specialist
Tyler Mitchell, Northeast Farm, ag specialist

Introduction

Corn and soybean planting is one of the most critical operations of the season. Operating the planter with the proper soil conditions for proper placement of the seed to obtain the correct seed-soil contact is important to optimize yields and reduce problems later in the season with plant and root growth. As corn and soybean seed prices continue to rise, and grain prices fall, it is important for farmers to find a population that maximizes both yield and profit. Planting too high of a corn population can result in increased barrenness and thus lower yields, but too low of a population also can result in lower yields. Past studies have indicated soybean yields are similar across a wide range of populations, but too low of a population can result in reduced yields and too high of a population can reduce profits. The objective of these trials was to investigate the effect of various planter operations on corn and soybean yield.

Materials and Methods

In 2017, eight trials investigated the effects of various aspects of corn planter operations on corn yield (Table 1), and six trials investigated the effects of various planter operations on soybean yield (Table 2). All trials were

conducted on-farm by farmer cooperators using the farmer's equipment. Strips were arranged in a randomized complete block design with at least three replications per treatment. Strip length and width varied from field-to-field depending on field and equipment size. All plots were machine harvested for grain yield.

Corn Trial 1 compared using precision planting E-set plates with John Deere 30 cell plates (Table 3). In Trial 2, the effect of planter speeds from 4 mph to 7.5 mph on corn yield was investigated. Trials 3, 4, 5, 6, 7, and 8 investigated the effect of two to three seeding rates on corn yield. Trial 5 also investigated using two different nitrogen (N) rates with the three different seeding rates. A preplant N rate of 170 lb N/acre with or without an additional rate of 50 lb N/acre side-dressed at V6 crop growth stage was investigated in Trial 5.

In soybean Trial 1, soybeans planted in late April were compared with soybeans planted in mid-May (Table 4). In Trial 2, the effect of row width on soybean yield was investigated. In Trials 3, 4, 5, and 6, the effect of seeding rate on soybean yield was investigated.

Results and Discussion

In corn Trial 1, corn planted using John Deere 30 cell plates yielded seven bushels/acre more than corn planted with precision planting E-set plates (Table 3). It is not known why this yield difference occurred. In Trial 2, there was no difference in corn yield with planting speeds ranging from 4 to 7.5 mph. In Trials 3, 4, and 7, there was no difference in corn yield with the three different seeding rates, with seeding rates varying from 28,000 to 39,000 seeds/acre. In Trial 5, there was no difference in corn yield with the three seeding rates from 34,500 to 38,000 seeds/acre and no difference

in corn yield with the two nitrogen fertilizer rates. In Trials 6 and 8, there was a higher yield with the lowest seeding rate of 34,000 seeds/acre compared with a seeding rate of 36,000-38,000 seeds/acre ($P \leq 0.07$). Based on these trials it is apparent seeding rates of 34,000 seeds/acre or less is adequate for maximum corn yields. Thus, there may be opportunities for some farmers to reduce their seeding rates, although results will likely vary from year to year.

In soybean Trial 1, there was no difference in yield between the soybeans planted in late April compared to soybeans planted in mid-May (Table 4). It is usually recommended to plant soybeans in late April to early May for maximum yields. In Trial 2, soybeans planted in 30-in. rows yielded four bushels/acre more than soybeans planted in 15-in. rows ($P = 0.07$). There was more white mold present in the soybeans planted in the 15-in. rows, which may have been responsible for the lower yield. Most studies have shown if there is a yield

advantage, it is usually soybean planted with the narrow rows that out-yield soybean in wider rows, unless white mold or other disease problems are in the field. In Trials 3, 5, and 6, there was no difference in soybean yield with the different seeding rates, with seeding rates varying from 125,000 seeds/acre to 180,000 seeds/acre. In Trial 4, the highest seeding rate of 150,000 seeds/acre resulted in a higher soybean yield than seeding rates of 90,000 to 130,000 seeds/acre ($P = 0.02$). It is usually recommended to seed about 140,000 seeds/acre in order to have a final plant stand of 100,000 plants/acre or more. The results of these trials support that recommendation.

NOTE: The results presented are from replicated demonstration trials. Statistics are used to detect differences at a location and should not be interpreted beyond the single location.

Table 1. Hybrid, row spacing, planting date, planting population, previous crop, and tillage practices in the 2017 planter trials on corn.

| Exp. no. | Trial | County | Hybrid | Row spacing (in.) | Planting date | Planting population (seeds/ac) | Previous crop | Tillage |
|----------|-------|------------|------------------|-------------------|---------------|--------------------------------|---------------|-----------------------------------|
| 170113 | 1 | Sioux | Pioneer PO589AMX | 30 | 4/24/17 | 34,000 | Soybean | Conventional |
| 170810 | 2 | Howard | Pfister 6081 | 30 | 5/17/17 | 36,000 | Corn | No-till |
| 170207 | 3 | Crawford | Curry 826-64AMX | 30 | 5/20/17 | 30,000, 34,000, 38,000 | Corn | Fall disk, spring field cultivate |
| 170301 | 4 | Monona | Wyffels 7456VT2 | 30 | 5/26/17 | 28,000, 32,000, 36,000 | Corn | No-till |
| 170401 | 5 | Hancock | Becks 5441SX RIB | 30 | 5/8/17 | 34,500, 36,500, 38,000 | Soybean | Conventional |
| 170706 | 6 | Washington | Dekalb DK6435 | 30 | 4/17/17 | 34,000, 36,000 | Soybean | No-till |
| 170812 | 7 | Howard | Pioneer PO574AM | 30 | 5/10/17 | 33,000, 36,000, 39,000 | Corn | No-till |
| 170823 | 8 | Bremer | New Tech 5F-504 | 30 | 5/10/17 | 34,000, 38,000 | Corn | No-till |

Table 2. Variety, row spacing, planting date, planting population, previous crop, and tillage practices in the 2017 planter trials on soybean.

| Exp. no. | Trial | County | Variety | Row spacing (in.) | Planting date | Planting population (seeds/ac) | Previous crop | Tillage |
|-----------------|--------------|---------------|-----------------|--------------------------|---------------------------|--|----------------------|-------------------------|
| 170119 | 1 | Sioux | Pioneer P18T85R | 30 | 4/24/17 and 5/15/17 | 140,000 | Corn | No-till |
| 170127 | 2 | Lyon | Becks 185R2 | 15 and 30 | 5/10/17 | 140,000 | Corn | Strip till |
| 170128 | 3 | Osceola | Asgrow A20XY | 30 | 5/13/17 | 125,000, 150,000, 175,000 | Corn | Conventional |
| 170129 | 4 | Sioux | Pioneer P24T56L | 30 | 5/11/17 | 90,000, 110,000, 130,000, 150,000 | Soybean | No-till |
| 170213 | 5 | Pocahontas | Syngenta S20T6 | 30 | 5/15/17 | 149,735, 179,742 | Corn | Chisel, field cultivate |
| 170408 | 6 | Franklin | Syngenta S26-P3 | 30 | 5/12/17 | 145,000 165,000 | Corn | Conventional |

Table 3. Yields for on-farm corn planter trials in 2017.

| Exp. no. | Trial | Treatment | Spring stand count (plants/ac) | Fall stand count (plants/ac) | Yield (bu/ac) ^a | P-value ^b |
|----------|-------|--|--------------------------------|------------------------------|----------------------------|----------------------|
| 170113 | 1 | Planted with Precision E-set plates | | | 261 a | 0.01 |
| | | Planted with John Deere 30 cell plates | | | 268 b | |
| 170810 | 2 | Planted at 4.0 miles/hour | | | 198 a | 0.57 |
| | | Planted at 5.0 miles/hour | | | 200 a | |
| | | Planted at 5.5 miles/hour | | | 203 a | |
| | | Planted at 6.0 miles/hour | | | 195 a | |
| | | Planted at 6.5 miles/hour | | | 187 a | |
| | | Planted at 7.0 miles/hour | | | 205 a | |
| | | Planted at 7.5 miles/hour | | | 205 a | |
| 170207 | 3 | Planted at 30,000 seeds/ac | 28,300 a | 27,000 a | 230 a | 0.19 |
| | | Planted at 34,000 seeds/ac | 31,000 b | 30,500 b | 236 a | |
| | | Planted at 38,000 seeds/ac | 35,300 c | 34,600 c | 236 a | |
| 170301 | 4 | Planted at 28,000 seeds/ac | 28,600 a | 28,000 a | 214 a | 0.17 |
| | | Planted at 32,000 seeds/ac | 32,400 b | 31,900 b | 225 a | |
| | | Planted at 36,000 seeds/ac | 36,900 c | 35,800 c | 234 a | |
| 170401 | 5 | Planted at 34,500 seeds/ac with 170 lb/ac N as UAN preplant | | | 298 a | 0.34 |
| | | Planted at 34,500 seeds/ac with 170 lb/ac N as UAN preplant plus 50 lb/ac N as UAN sidedress at V6 | | | 294 a | |
| | | Planted at 36,000 seeds/ac with 170 lb/ac N as UAN preplant | | | 300 a | |
| | | Planted at 36,000 seeds/ac with 170 lb/ac N as UAN preplant plus 50 lb/ac N as UAN sidedress at V6 | | | 299 a | |
| | | Planted at 38,000 seeds/ac with 170 lb/ac N as UAN preplant | | | 303 a | |
| | | Planted at 38,000 seeds/ac with 170 lb/ac N as UAN preplant plus 50 lb/ac N as UAN sidedress at V6 | | | 306 a | |
| 170706 | 6 | Planted at 34,000 seeds/ac | | | 242 a | |
| | | Planted at 36,000 seeds/ac | | | 234 b | |
| 170812 | 7 | Planted at 33,000 seeds/ac | 31,300 a | | 226 a | 0.39 |
| | | Planted at 36,000 seeds/ac | 34,300 ab | | 232 a | |
| | | Planted at 39,000 seeds/ac | 36,800 b | | 220 a | |
| 170823 | 8 | Planted at 34,000 seeds/ac | | 32,000 a | 214 a | 0.07 |
| | | Planted at 38,000 seeds/ac | | 34,000 a | 209 a | |

^aValues denoted with the same letter within a trial are not statistically different at the significance level of 0.05.

^bP-value = the calculated probability that the difference in yields can be attributed to the treatments and not other factors. For example, if a trial has a P-value of 0.10, then we are 90 percent confident the yield differences are in response to treatments. For P = 0.05, we would be 95 percent confident.

Table 4. Yields for on-farm soybean planter trials in 2017.

| Exp. no. | Trial | Treatment | Fall stand count (plants/ac) | Yield (bu/ac) ^a | P-value (yield) ^b |
|----------|-------|-----------------------------|------------------------------|----------------------------|------------------------------|
| 170119 | 1 | Planted 4/24/17 | | 70 a | 0.20 |
| | | Planted 5/15/17 | | 69 a | |
| 170127 | 2 | Planted with 15-in. rows | | 62 a | 0.07 |
| | | Planted with 30-in. rows | | 66 a | |
| 170128 | 3 | Planted 125,000 seeds/ac | | 70 a | 0.30 |
| | | Planted 150,000 seeds/ac | | 70 a | |
| | | Planted 175,000 seeds/ac | | 70 a | |
| 170129 | 4 | Planted 90,000 seeds/ac | 86,000 a | 74 a | 0.04 |
| | | Planted 110,000 seeds/ac | 102,000 ab | 74 a | |
| | | Planted 130,000 seeds/ac | 119,000 bc | 74 a | |
| | | Planted 150,000 seeds/ac | 143,000 c | 76 b | |
| 170213 | 5 | Planted at 150,000 seeds/ac | 120,900 a | 59 a | 0.86 |
| | | Planted at 180,000 seeds/ac | 141,200 b | 60 a | |
| 170408 | 6 | Planted at 145,000 seeds/ac | | 55 a | 0.74 |
| | | Planted at 165,000 seeds/ac | | 55 a | |

^aValues denoted with the same letter within a trial are not statistically different at the significance level of 0.05.

^bP-value = the calculated probability that the difference in yields can be attributed to the treatments and not other factors. For example, if a trial has a P-value of 0.10, then we are 90 percent confident the yield differences are in response to treatments. For P = 0.05, we would be 95 percent confident.