

## Fairway Height Bentgrass Cultivar Trial

### RFR-A2014

Nick Christians, university professor  
Adam Thoms, assistant professor  
Ben Pease, research scientist  
Department of Horticulture

### Introduction

This is the final year of data collection from the fairway height bentgrass trial established in the fall of 2014. The site was killed in August to make way for a new trial.

### Materials and Methods

The area was maintained at a 0.4 in. mowing height. This is a National Turfgrass Evaluation (NTEP) trial and is being conducted at several research stations in the United States. It contains 17 of the newest cultivars and a number of experimental cultivars. The cultivars are maintained with 4 lb of N/1,000 ft<sup>2</sup> per growing season. Fungicides are used as needed in a preventive program. Herbicides and insecticides also are applied as needed. Irrigation was applied as needed to prevent moisture stress.

### Results and Discussion

The visual quality was evaluated monthly in 2020 from April through August (Table 1). The values listed under each month in Table 1 are the averages of visual quality ratings made on three replicated plots. Visual quality was based on a scale of 9 to 1: 9 = best quality, 6 = lowest acceptable quality, and 1 = worst quality. The annual average of the monthly data for each cultivar was calculated and is listed in the last column (Mean). Data for Brown Patch also were collected in July. The scale is 1 = no disease and 9 = worst. The last row states the LSD (least significant difference), which is a statistical measurement of how widely the data in each column must vary before they are considered to be different from one another. Spring greenup was rated in April using a 9 to 1 scale with 9 = dark and 1 = light green. NS at the bottom of the column means there were no significant differences in the data in that column.

### Acknowledgements

Appreciation is extended to the NTEP organization for funding to conduct this project.

**Table 1. Data for the fairway height creeping bentgrass trial-2020.**

Cultivar	Quality <sup>a</sup>						BrownP <sup>b</sup>	SpringG <sup>c</sup>
	April	May	June	July	Aug	Mean		
DLFPS-AT/3026	6	6	6	7	5	6	1	6
7	6	8	8	6	4	6	2	6
Penncross	6	6	6	7	6	6	1	6
Crystal Blue Links	6	6	7	7	8	7	1	5
Greentime	5	7	6	8	5	6	2	6
L-93XD	6	6	7	7	4	6	1	5
Armor	5	6	6	7	5	6	2	5
Kingdom	6	5	6	7	4	6	1	5
Nightlife	5	6	7	6	6	6	1	4
V-8	6	7	7	7	6	7	1	6
DC-1	5	7	6	7	5	6	1	6
PPG-AT 104	5	7	6	7	4	6	1	5
H10G-OP	6	7	7	6	6	7	3	5
Shark	5	7	6	7	7	6	1	6
Barracuda	7	8	7	7	6	7	1	7
PST-ORBS	6	6	7	6	6	6	2	6
PST-DCV6	6	6	7	7	8	7	1	6
LSD 0.05	1	NS	1	NS	3	1	NS	NS

<sup>a</sup>Quality rated on a 1 to 9 scale with 9 = best quality, 1 = worst quality, and 6 = lowest acceptable quality.

<sup>b</sup>BrownP is brown patch: 1 = no disease and 9 = worst.

<sup>c</sup>Spring greenup rated on a 1 to 9 scale with 9 = dark, 1 = light green.