

2007

## WinterTriticale Variety Test

Ronald Skrdla  
*Iowa State University*

Jean-Luc Jannink  
*Iowa State University*

Follow this and additional works at: [http://lib.dr.iastate.edu/farms\\_reports](http://lib.dr.iastate.edu/farms_reports)



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), and the [Agronomy and Crop Sciences Commons](#)

---

### Recommended Citation

Skrdla, Ronald and Jannink, Jean-Luc, "WinterTriticale Variety Test" (2007). *Iowa State Research Farm Progress Reports*. 962.  
[http://lib.dr.iastate.edu/farms\\_reports/962](http://lib.dr.iastate.edu/farms_reports/962)

This report is brought to you for free and open access by Iowa State University Digital Repository. It has been accepted for inclusion in Iowa State Research Farm Progress Reports by an authorized administrator of Iowa State University Digital Repository. For more information, please contact [digirep@iastate.edu](mailto:digirep@iastate.edu).

---

# WinterTriticale Variety Test

**Abstract**

Twelve varieties were included in the 2006 winter triticale variety test at Sutherland. Each variety was sown in three different plots to average the effects of soil variability. The varieties were planted October 3, 2005 at a rate of 1½ bushels/acre. All winter triticale plots were harvested on July 18.

**Keywords**

Agronomy

**Disciplines**

Agricultural Science | Agriculture | Agronomy and Crop Sciences

## Winter Triticale Variety Test

Ron Skrdla, ag research specialist  
Jean-Luc Jannink, assistant professor  
Department of Agronomy

### Materials and Methods

Twelve varieties were included in the 2006 winter triticale variety test at Sutherland. Each variety was sown in three different plots to average the effects of soil variability. The varieties were planted October 3, 2005 at a rate of 1½ bushels/acre. All winter triticale plots were harvested on July 18.

### Results and Discussion

Winter triticale yields averaged 94.9 bushels/acre in 2006, which is 27.3 bushels/acre more than the statewide long-term average (Table 1). NE426GT was the highest yielding line based on the long-term average while Décor had the highest test weight across all locations for the lines that were tested in 2006.

Additional information on barley variety tests in the state can be found in the publication, “Iowa Crop Performance Tests—Winter Wheat and Winter Triticale, 2006,” which is available from county extension offices (Pm-1645) and at [www.public.iastate.edu/~jjannink/](http://www.public.iastate.edu/~jjannink/).

**Table 1. Performance of winter triticale varieties tested at Sutherland in 2006.**

Variety	Yield				
	2006	Long-term avg.	Test weight <sup>1</sup> (lb/bu)	Heading date <sup>2</sup> (May)	Plant height <sup>3</sup> (in.)
Alzo	82.8	64.0	51.1	26.0	37.9
Arapahoe <sup>4</sup>	94.8	57.7	58.2	26.0	33.8
Danko Presto	92.3	67.4	54.4	22.4	40.3
Décor	96.7	69.3	56.9	20.4	38.8
Kitaro	103.7	70.9	55.8	23.0	36.1
Lamberto	107.7	73.9	53.8	25.0	38.1
NE422T	88.2	64.7	54.1	29.3	48.2
NE426GT	111.0	78.3	53.8	22.7	39.3
Pika	83.6	54.9	54.7	34.8	51.3
Sorento	99.3	73.9	52.8	24.7	38.4
Trical 336	86.6	61.6	53.8	23.0	40.4
Trical 815	104.1	72.9	53.6	23.7	40.3
Vero	83.0	70.4	53.2	24.0	41.2
Average	94.9	67.6	54.3	25.0	40.3
LSD(0.05) <sup>5</sup>	11.3	10.5	2.5	3.2	4.3

<sup>1</sup>Test weight—average from three sites.

<sup>2</sup>Data collected at Ames only recorded as date after May 1.

<sup>3</sup>Height—measured at Ames.

<sup>4</sup>Arapahoe—winter wheat variety used as a check.

<sup>5</sup>LSD=least significant difference. When entries differ by an amount equal to one LSD or more, they are considered to be in different classes with 95% certainty.