

2007

# Evaluation of Organic Soybean Rust Treatments for Organic Production

Kathleen Delate

*Iowa State University*, [kdelate@iastate.edu](mailto:kdelate@iastate.edu)

Andrea McKern

*Iowa State University*

Robert Burcham

*Iowa State University*

Follow this and additional works at: [http://lib.dr.iastate.edu/farms\\_reports](http://lib.dr.iastate.edu/farms_reports)



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), [Agronomy and Crop Sciences Commons](#), and the [Horticulture Commons](#)

---

## Recommended Citation

Delate, Kathleen; McKern, Andrea; and Burcham, Robert, "Evaluation of Organic Soybean Rust Treatments for Organic Production" (2007). *Iowa State Research Farm Progress Reports*. 832.

[http://lib.dr.iastate.edu/farms\\_reports/832](http://lib.dr.iastate.edu/farms_reports/832)

This report is brought to you for free and open access by Iowa State University Digital Repository. It has been accepted for inclusion in Iowa State Research Farm Progress Reports by an authorized administrator of Iowa State University Digital Repository. For more information, please contact [digirep@iastate.edu](mailto:digirep@iastate.edu).

---

# Evaluation of Organic Soybean Rust Treatments for Organic Production

## **Abstract**

Asian soybean rust has the potential to be the single most important impediment to economical organic production. The economic impact in organic systems range from \$30 to \$120 million in yield loss. The fungus (*Phakopsora pachyrhiz*) survives year-round in warm areas, such as the southern United States. During a growing season, the pathogen is disseminated by northward seasonal wind.

## **Keywords**

Horticulture, Agronomy

## **Disciplines**

Agricultural Science | Agriculture | Agronomy and Crop Sciences | Horticulture

# Evaluation of Organic Soybean Rust Treatments for Organic Production

Kathleen Delate, associate professor  
Andrea McKern, research assistant  
Departments of Horticulture and Agronomy  
Bob Burcham, ag specialist

## Introduction

Asian soybean rust has the potential to be the single most important impediment to economical organic production. The economic impact in organic systems range from \$30 to \$120 million in yield loss. The fungus (*Phakopsora pachyrhiz*) survives year-round in warm areas, such as the southern United States. During a growing season, the pathogen is disseminated by northward seasonal wind.

Dry conditions across the Midwest in 2006 kept soybean rust out of Iowa, but soybean rust was found as far north as Indiana and has been shown to overwinter in Florida. Iowa State University has been awarded a grant from USDA to investigate “Strategies for Management of Asian Soybean Rust in Organic Systems,” which includes treatments allowable under certified organic conditions. Trials were established in 2005 in an area of Florida where rust is present to examine the effect of these treatments. Concurrently, trials have been established in Iowa, Pennsylvania, and Michigan to examine yield effects of these treatments under non-rust conditions.

## Materials and Methods

In the soybean rust treatment trial, Pioneer 9305 soybeans were planted at the Neely-Kinyon Farm on May 19, 2006, at 200,000 seeds/acre. Plots measuring 10 ft × 20 ft were laid out in a randomized complete design. There were four replications of the following treatments: MicroAF™ at 1 gal/acre, Sonata® at 1 gal/acre, and AgriCoat Natural IV™ at 12 oz/acre. All treatments were compared with a control. Treatments were applied on July 14, 2006, at the

R-1 stage. Leaves were inspected for disease on July 13, 26, August 11, 29, and September 13, by randomly selecting one leaf from the top, middle, and bottom sections of four plants per plot. Plots were maintained with one rotary hoeing on June 1, one cultivation on June 22, and walking on August 3, 2006. Soybeans were harvested on October 31. The percentage of stained soybeans was determined by counting the number of stained soybeans in a 200-gram sample that was randomly collected from the harvest of each plot.

## Results and Discussion

Soybean rust was not observed in the trial plots in 2006. The treatments had no effect on yield, soybean diseases, or grain quality (Table 1). Other diseases observed in 2007 include Frog Eye Leaf Spot, Cercospora Leaf Blight, Bacterial Pustule, Bacterial Blight, Brown Spot, and Downy Mildew. The most prevalent diseases were Frog Eye Leaf Spot and Downy Mildew (Figure 1). The highest number of infected leaves from Frog Eye Leaf Spot and Downy Mildew occurred on August 29 and August 11, respectively. We will continue to test products in 2007 and monitor treatment effects on other soybean diseases, in the event of no rust appearing.

## Acknowledgments


We would like to thank the Leopold Center for Sustainable Agriculture, the Wallace Foundation, Greg Lilly, David Rosmann, Magda Marin Beltran, Kristin Brown, Daniel Rosmann, Mark Rosmann, Rita Tisinger, Christina Erickson, and Francisco Viteri for their help. We also thank Eric Gross of Pioneer Hi-Bred International, TerraMax, Inc., Cottage Grove, MN, AgraQuest, Inc., Davis, CA, and AgriCoat LLC, Soledad, CA; and Charles Hurburgh and Glen Rippke of the ISU Grain Quality Lab for grain analysis.

**Table 1. Yield and grain quality in soybean rust treatment trial, Neely-Kinyon, 2006.**

Treatment	Yield (bu/acre)	Protein (%)	Oil (%)	Fiber (%)	Carbohydrates (%)	Moisture (%)
Control	43.88	37.64	17.83	4.70	21.84	10.93
Sonata®	46.09	37.39	17.96	4.70	21.95	10.84
AgriCoat Natural IV™	49.78	37.43	17.80	4.70	22.08	10.90
MicroAF™	44.25	37.70	17.63	4.70	21.98	10.83
LSD 0.05	NS	NS	NS	NS	NS	NS

**Table 2. Yield and grain quality in soybean rust treatment trial, Neely-Kinyon, 2006.**

Treatment	Stained soybeans (%)	Disease presence average over the growing season (%)					
		Frogeye leaf spot	Cercospora leaf blight	Bacterial pustule	Bacterial blight	Brown spot	Downy mildew
Control	5.43	22.6	1.3	0.4	1.3	0.9	12.2
Sonata®	3.84	18.3	2.6	1.7	2.2	0.0	12.7
AgriCoat Natural IV™	5.58	21.7	3.4	1.7	1.7	0.0	15.3
MicroAF™	4.48	19.2	2.1	0.9	1.3	0.0	14.1
LSD 0.05	NS	NS	NS	NS	NS	NS	NS

**Figure 1. Soybean disease occurrence in soybean rust treatment trial, Neely-Kinyon, 2006.**