

2013

Optimum Soybean Planting Date

Wayne B. Roush

Iowa State University, wroush@iastate.edu

Follow this and additional works at: http://lib.dr.iastate.edu/farms_reports



Part of the [Agricultural Science Commons](#), and the [Agriculture Commons](#)

Recommended Citation

Roush, Wayne B., "Optimum Soybean Planting Date" (2013). *Iowa State Research Farm Progress Reports*. 1995.
http://lib.dr.iastate.edu/farms_reports/1995

This report is brought to you for free and open access by Iowa State University Digital Repository. It has been accepted for inclusion in Iowa State Research Farm Progress Reports by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Optimum Soybean Planting Date

Abstract

Past research by Iowa State University has shown that the optimum planting date for soybeans, assuming favorable soil conditions, is the first week in May for the northern third of Iowa. The optimum date for the southern two thirds of Iowa is the last week of April. Given that rapidly changing soybean genetics have shown improvements in both yield and disease resistance, these two trials were designed to demonstrate the planting recommendation under local conditions.

Disciplines

Agricultural Science | Agriculture

Optimum Soybean Planting Date

RFR-A1268

Wayne Roush, farm superintendent

Introduction

Past research by Iowa State University has shown that the optimum planting date for soybeans, assuming favorable soil conditions, is the first week in May for the northern third of Iowa. The optimum date for the southern two thirds of Iowa is the last week of April. Given that rapidly changing soybean genetics have shown improvements in both yield and disease resistance, these two trials were designed to demonstrate the planting recommendation under local conditions.

Materials and Methods

The two trials were on a Monona silt loam soil type with the majority having a 2 to 5 percent slope. The site was located in Monona County with the last week in April as the recommended planting date. The trials were replicated four times with two different planting date treatments. Planting dates were April 30 and May 14. Plot size was 20 ft wide by approximately 540 ft long. The trials had no fall or spring tillage and were no-till planted into standing corn residue in 30-in. rows. Seed drop for both trials was at 138,898/acre.

Dry fertilizer was broadcast spread in the spring as a mixture of 11-52-0 and 0-0-60 yielding an analysis of 17-80-80/acre. One pre-plant burndown and one post-emerge application of Roundup was used. The burndown also contained 2 oz of Sencor DF and 0.8 oz of Brawl II. Weed control was rated as excellent.

The two trials were located in the same field with all planting, fertilizing, spraying, and harvesting dates and rates the same. The same

seed variety (Renze 2889RR) was used in both trials. However, Trial 1 had a CruiserMax seed treatment and Trial 2 did not have any seed treatment.

Results and Discussion

Tables 1 and 2 show that there was a yield benefit for early planting in both trials in 2012. These yields reinforce the ISU recommendation that for the southern two thirds of Iowa, under favorable soil conditions, the optimal soybean planting date is the last week of April. It should be noted that planting soybeans too early also has some risks. Early planted soybeans may have more seedling diseases, may be at greater risk of sudden death syndrome, and could be damaged if a late spring frost occurs.

Acknowledgements

Appreciation is extended to Dennis Boyle of Renze Seeds for providing the seed used in this trial. Additional thanks to Brad Hanson for harvesting the plot.

Table 1. Treated seed soybean planting date yields.

Treatment	Yield (bu/acre)
April 30	65.0
May 14	54.4*

* = statistical difference at a P value of P = 0.09.

Table 2. Non-treated seed soybean planting date yields.

Treatment	Yield (bu/acre)
April 30	60.1
May 14	51.9*

* = statistical difference at a P value of P = 0.16.