

STERKIANA

NUMBER 62

COLUMBUS, OHIO

JUNE 1976

CONTENTS	PAGE
ROY STANLEY RIGGLE -- QUANTITATIVE EXAMINATION OF GASTROPOD AND SOIL RELATIONSHIPS IN AN OAK-HICKORY FOREST IN THE LOWER ILLINOIS VALLEY REGION	1
RALPH W. DEXTER -- POSTSCRIPT ON MUSSEL VS. MUSCLE IN RELATION TO BIVALVES	18
RAYMOND W. NECK -- MICRO-DISTRIBUTION OF LAND SNAILS IN AN ARTIFICIAL TALUS SLOPE	19
BRANLEY A. BRANSON -- <i>TESTACELLA HALIOTIDEA</i> (DRAPARNAUD) FREE LIVING IN PENNSYLVANIA	20
REPRINTS OF RARE PAPERS ON MOLLUSCA: MÜLLER, 1774, VERMIUM TERRESTRIUM ET FLUVIATILIMUM, SEU ANIMALIUM INFUSORIORUM, HELMINTHICORUM, ET TESTACEORUM, NON MARINORUM SUC-CINCTA HISTORIA HAVNIAE ET LIPSIAE, APUD HEINECK ET FABER. (FOREWORD)	20

.....

EDITORIAL BOARD

HENRY VAN DER SCHALIE
UNIVERSITY OF MICHIGAN
ANN ARBOR, MICHIGAN

WILLIAM J. WAYNE
UNIVERSITY OF NEBRASKA
LINCOLN, NEBRASKA

DAVID H. STANSBERY
OHIO STATE UNIVERSITY
COLUMBUS, OHIO

AURELE LA ROCQUE
OHIO STATE UNIVERSITY
COLUMBUS, OHIO

EDITOR

AURELE LA ROCQUE
102 W. BEAUMONT ROAD
COLUMBUS, OHIO 43214

NOTICE TO SUBSCRIBERS

RATE CHANGES

Since publication of the first number of STERKIANA in 1958, the price of a single number has been 50¢ prepaid. In the last 13 years, the price of paper has doubled and postage rates have almost tripled. It has become necessary to increase the price of subscriptions as follows:

Yearly subscriptions, 4 numbers paid in advance	\$ 2.20
Invoicing charge when required	1.00
Single back numbers (except No. 50) mailed individually	0.60
Back numbers, in lots of 10 or more, per number, postpaid	0.55
Double number (No. 50) mailed individually	1.10

*Reprints of articles in STERKIANA
are available at the following rates:*

*One sheet (1 or 2 pages): \$1.00 per
hundred copies, plus postage.
This price does not include collat-
ing or special covers.*

Authors should note that illustrations requiring commercial plate making must be charged to the author.

Subscribers should note that normally, the last number paid for has a notice of expiry of subscription on the last page. There is no invoicing charge if this notice is sent with remittance for the next four numbers.

QUANTITATIVE EXAMINATION OF GASTROPOD AND SOIL
RELATIONSHIPS IN AN OAK-HICKORY FOREST
IN THE LOWER ILLINOIS VALLEY REGION

ROY STANLEY RIGGLE

UNIVERSITY OF WISCONSIN, MADISON, WISCONSIN

ABSTRACT

Malacology can contribute significantly to the reconstruction of the local ecology at archaeological sites. Detailed reconstructions, however, are hampered by the paucity of molluscan ecological information, particularly at the local level.

Seven soil variables and their effects on the distribution of thirteen snail species were examined for part of a mesic oak-hickory stand in the lower Illinois Valley region. Generalized relationships between soil variables and snails were apparent. The number of species, and less so the number of specimens, tends to increase as the sample area increases. Traditional methods of graphic analysis and bivariate statistics did not provide optimal information on underlying relationships between the data sets. Few significant bivariate correlations were obtained between snail species and soil variables.

There was generally a nonlinear or negative cor-

relation between snail species and soil pH. No significant bivariate correlations were found between snail species and organic matter, available phosphorus, acid soluble phosphorus or exchangeable magnesium. A high positive correlation existed between calcium and three of thirteen species but was significant only between *Strobilops labyrinthica* (Say) and calcium. A generally positive but not significant correlation existed between three species widely distributed over North America, and all measured soil parameters except exchangeable potassium.

Factor analysis delineated three underlying patterns in the variability and covariance of the data and provided a basis for inferring the secondary influence of soil factors on the distribution of snails. Variables which covary in each eigenvector were usually snail species rather than snail and soil variables. Habitat groups of snails were defined on the basis of species with a similar pattern of covariation.

INTRODUCTION

The environment of prehistoric village and mortuary sites has become a subject of increasingly popular interest among archeologists for several decades. This interest is warranted due to the current orientation of archeological research--an evaluation of man's interaction in an ecosystem.

Terrestrial mollusks, because of their sensitivity to local habitat conditions, are useful in the reconstruction of past environments. Interpretation of gastropod assemblages from archeological contexts rests partly on knowledge of presently living species. Analysis of archeological assemblages

requires intensive and extensive sampling at the local level of the site. There are few analyses of living snails, particularly at the local level, which are sufficiently detailed to use as analogs in the explanatory and interpretative framework for assemblages from prehistoric sites.

The present study is a quantitative examination of relationships between gastropods and soil factors on the surface of a burial mound in a mesic oak-hickory forest. It deals intensively with micro-habitat variation in soils and snails and quantitatively measures the covariation of each.

ACKNOWLEDGEMENTS

The author is indebted to Dr. Theodore R. Peck (Department of Agronomy, University of Illinois, Urbana) who provided detailed soil analyses. Without Dr. Peck's cooperation the analysis could only have been a very abbreviated form of the one presented here. In addition, Dr. Wayne Wendland (Department of Meteorology, University of Wisconsin, Madison) offered very helpful suggestions on eigen-vector analysis.

DESCRIPTION OF PROJECT AREA

The Mueller-Ringhausen forest, on the property of Mr. George Mueller, is located (Fig. 1) about 2.5 miles north of the town of Hamburg, Illinois, about one mile east of the east bank of the Mississippi River (northeast corner of northeast quarter of southwest quarter of Section 14 Township 9 South, Range 3 West).

Much of the area of upland slopes of interior Calhoun County supports oak-hickory forests. The long-standing deciduous forests, in addition to climate, are primarily responsible for thick well developed Alfisol soils common in the county.

The collecting area is located on a northwest-southeast oriented ridge which drops with a gentle slope to the north. The area is adjacent to a broad plateau around which are numerous dissecting valleys. A group of four Late Woodland burial mounds is located on one of the ridges. The gastropod assemblage discussed below was recovered from the surface of Mound 2. The surface of this mound was selected for the study because of its degree of preservation and the presence of what appeared to be a nearly homogeneous microhabitat over the area of the mound. The vegetation on Mounds 3 and 4, located near a fence line bordering an open pasture, had been greatly altered into edge area habitats and Mound 1, to the north, was the target of recent disturbances by collectors.

FORMULATION OF HYPOTHESES

The effect of soil factors on the distribution and abundance of terrestrial gastropods has been previously investigated on a regional scale in three counties in eastern Virginia (Burch 1955, 1956). Contrary to Boycott (1929) and Oughton (1948) Burch concluded that organic matter did influence the presence of land snails, thus agreeing with Shimek (1930) that organic matter as a potential food is a major limiting factor in the distribution of land snails.

Burch also investigated several other soil parameters. He concluded that pH did not affect the distribution of land snails. Research by other investigators (Strandine 1941) has not established a clear systematic relationship between this parameter and terrestrial gastropods. This variable is presently considered to be of indirect or negligible importance to the distribution of snails.




FIGURE 1. Topographic map showing position of samples of the stratified systematic unaligned sample of the surface of Mound 2 of the Mueller-Ringhausen mound group. Location of samples is approximate with respect to topography since they were obtained by measuring along ground surface. Dashed line is approximate perimeter of mound fill.

With respect to other soil factors snail frequencies were found to increase as the concentration of calcium (CaO), magnesium (MgO) and potassium (K_2O) in the soil increased. Potassium in addition to phosphorus and pH were concluded to have an indirect effect on the distribution of land snails (Burch 1955, 1956). Other investigators (e. g. Atkins 1966) demonstrated a positive relationship between an increase in calcium carbonate ($CaCO_3$) and an increase in frequency and species per unit area in alkaline soils. In all cases although the variables are quantified by the respective investigator an actual measure of correlation is not provided.

The present study attempts to determine the quantitative degree of covariance between gastropods and soil parameters similar to those used by Burch, on a local level. The soil variables investigated were organic matter content, available phosphorus, acid soluble phosphorus, exchangeable potassium,

exchangeable calcium, exchangeable magnesium and soil pH. All of the soil parameters measured are interrelated, and are a product of the interactive system of soil and the non-soil environment, e. g. vegetation and climate. The latter variables are, however, important in the development of a soil but for the purposes of this analysis are treated separately.

Previous studies (cf. Burch 1955, 1956) have not included the quantitative explanation of local variability in soil parameters as they relate to gastropods, nor the investigation of alternate forms in phosphorus can be extracted. Burch's study, limited to a region of eastern Virginia, gives important clues to the ecological relationships of gastropods and soils. In each location for which soil data were measured, however, the sample area at each site was not subdivided. This produces an average of soil conditions which may or may not show the range of variability in each area. Secondly, the distribution of species may change markedly within a few feet.

Numerous investigators have contributed to a large body of literature on the occurrence and abundance of species in forests of varying vegetational composition (e. g. Van Cleave 1951; Shimek 1930). The delineation of a complex of species characteristic of forests has proceeded on a qualitative basis for many years. The composition of the habitat group should be expected to change between regions due to the effect of changes in marked interregional climatic patterns. Within the same region species comprising a habitat group likely change on a continuum responding to the primary factors of vegetation (cover) and moisture. Plant ecologists have given ample demonstration of a vegetational continuum rather than the existence of discrete vegetational units (cf. Curtis 1959). Significant positive correlation between species diversity and moisture regime and between snail diversity and the number of dominant tree species has been documented for eight habitats in the Great Smoky Mountains (Getz 1974).

Quantitative definition of habitat groups as used in this study is based on the division of an assemblage into groups of species with the greatest similarity of covariance. Within each habitat spatial variation in the distribution of species and frequencies can be used to detect minor changes in a contemporaneous population. Some of the changes can be expected to be related to the autecology of the species present in the area. By examining many sites in a given region the soil and gastropod variables which show systematic variation take on regional significance.

The present analysis examines the relationship between gastropods and soils in a part of one stand of oak-hickory forest. If the relationship between gastropods and a complex of soil variables can be quantitatively defined on a local level gastropod and soil analyses of archeological sites can be used to provide substantial reconstruction of local environmental conditions. It is clear that the soil factors (with the probable exception of calcium)

affecting the spatial distribution of gastropods are minor compared to the effect of plant cover and moisture. Soil factors are interrelated and reflect plant conditions. Some soil parameters change markedly during the depositional history of an archeological site. Therefore it is important to focus on the more stable variables of soil which undergo minor changes through time; e. g., phosphorus, and bulk density. Due to nutrient pumping by plants (Odum 1971) even these variables may change radically unless deposition on the surface is rapid. If gastropod distribution was affected directly by soil factors interpretation of archeological assemblages would be inextricably complex due to the nearly constant change in the soil environment.

Burial mounds provide a situation where buried humus zones are frequently encountered. Rapid addition of fill above humus layers effectively seals off the underlying strata and their contained flora and fauna. Alkaline soil environments are favorable for the preservation of gastropod shells and in some cases the spatial relationships of the assemblage may be preserved (death assemblage) with minor modifications. In such cases comparison of the archeological assemblage with assemblages recovered from known ecological situations may reveal evidence of spatial distribution of habitats over the area of the buried surface of the mound, or the original surface on which the mound was erected. These data can reveal the record of short term shifts in ecotones provided the mounds are sufficiently dense over an area and can be arranged chronologically.

SUMMARY OF HYPOTHESES

The surface gastropod assemblage of a part of the Mueller-Ringhausen forest was used to evaluate the following hypotheses.

1. The habitat group characteristic of the part of the oak-hickory forest sampled would be similar to the complex of species most common in other areas of oak-hickory forest. In this case however the delineation of the habitat group is largely quantitative. Due to its vegetational structure this stand may be expected to contain the species common in a qualitatively defined habitat group but in different proportions in response to local conditions.

2. The spatial relationships of gastropod species can be expressed quantitatively by analysis of their frequencies per unit area and summarized by their patterns of covariance. Generally the greater the frequency and relative diversity per unit area the greater will be the amount of contained information.

3. Minor changes or variation in the distribution of snails and associated soil data reveal minor changes in a relatively homogeneous macrohabitat.

4. Soil pH should have little effect on the distribution of snails in this particular macrohabitat. Organic matter, and secondarily calcium, magnesium, and potassium should generally increase as snail frequencies increase.

Each of these variables should covary with gastropod species which show positive response to increases in snail parameters.

FIELD AND LABORATORY METHODS AND MATERIALS

The strategy employed during the design of a sampling program for the surface of Mound 2 was to obtain samples from points over the entire extent of the mound surface. This necessitated a design which would insure obtaining samples from the highest point of the mound where vegetation was less dense, as well as on the slopes and edge of the mound which supported a thick layer of decomposing litter. The final choice of a design was the stratified systematic unaligned sample (Haggett 1965). The procedure offers the advantages of randomization, stratification and systematic sampling and avoids the alignment of sample points and thus the possibility of error due to periodicities in the phenomena being sampled (Haggett 1965: 196).

The position of samples over the surface of the mound is indicated in Figure 1. A 48-foot square was sufficient to cover the mound and some overlap in the corners off the circular mound was thus obtained. The square was divided into four quadrants and four samples were selected at random in each 24-foot square quadrant. In the field the samples were obtained by establishing a temporary datum point on the highest part of the approximate center of the mound. The sampling grid was oriented toward magnetic north. Tapes were then used to measure along ground distance along the four coordinate lines and out to a sample point at right angles to the coordinate lines. The measurements were taken along ground distance to the northwest corner of the one-foot square sample. Although the slope is relatively constant the snail sampling grid is not precisely related to the archeological grid by which the mound was subsequently excavated.

By shifting the points of origin of the X and Y axes of the sampling grid (cf. Haggett 1965: 196) five series of random samples were established. The first series contained 16 samples; the remaining four contained nine samples each. All but the fifth series of samples were collected. The first series, i.e. 16 samples, is reported in the present study.

Each sample consisted of two parts. All of the surface litter in one square foot down to the interface of the O2 and A1 soil horizons was placed in a plastic bag. A soil sample one inch deep and one foot square was placed in a separate plastic bag. The litter samples were subsequently soaked in a solution of sodium bicarbonate and water in screen-covered buckets before being washed in number 40-mesh screens and air-dried, followed by sorting. Sample processing and shell identification methods are discussed at length elsewhere (Riggle 1975; Jaehrig 1971).

The soil samples initially included all soil in one square foot for a depth of one inch below the surface litter. Subsequently each soil sample was

thoroughly mixed and approximately 300 milliliters was placed into heavy polyethylene bags and forwarded for analysis to Dr. Theodore R. Peck (Department of Agronomy, University of Illinois, Urbana).

Soil pH was determined by the paste method utilizing distilled water and a one-half hour equilibrium interval before the reading was obtained with a Beckman Zeromatic pH meter.

Organic matter was determined by the Walkly-Black procedure (Black 1965: 1372) and the results read colorimetrically.

Bray's P1 (Bray and Kurtz 1945) method was used to measure available phosphorus. Results were read colorimetrically. The extractant used was 0.03 normal ammonium fluoride (NH₄F) in 0.025 normal HCl. Acid soluble phosphorus was determined by Bray's P2 method using 0.03 normal NH₄F in 0.1 normal HCl as the extractant. The P1 method extracts phosphorus associated primarily with aluminum. Aluminum, in addition to some calcium phosphate, is extracted by the P2 method. The ratio of acid soluble to available phosphorus should be low in acid soils such as those present in the Mueller-Ringhausen forest.

Exchangeable potassium was extracted with 1 normal ammonium acetate at pH 7.0 and the result determined by flame photometry from a one to ten ratio of soil to extractant.

Exchangeable calcium and exchangeable magnesium were extracted using the method applied to the samples for extracting exchangeable potassium but a one to twenty ratio of soil to extractant was employed. The results were determined with atomic absorption spectrophotometry.

Botanical data collected during the field work were used to evaluate the relative amount each tree species contributed to the vegetation in the sample area. Although more rigorous procedures of establishing the characteristics of a stand are available (e. g. Curtis 1959) time did not permit exhaustive study of the vegetation.

Trees over three inches in diameter breast high were identified and counted. The trees on the surface of Mound 2 were predominantly white oak (*Quercus alba*), shagbark hickory (*Carya ovata*) and flowering dogwood (*Cornus florida*). Flowering dogwood and white oak trees were most common on the toe of the side slopes of the mound. One flowering dogwood and two white oak trees were located on the summit of the mound. Redbud (*Cercis canadensis*), dogwood (*Cornus* sp.) and white oak dominate the trees around the mound. Redbud and dogwood saplings (less than one-half inch in diameter) were sparsely distributed over the surface of the mound.

ANALYSIS AND DISCUSSION

Phenology and Taxonomy

Two problems which may have had some small ef-

fect on the quantitative analysis of the assemblage which follows this section are the phenology of certain snail species and the taxonomic assessment of one of the species.

The acidity of the soil samples obviates the possibility that the shells recovered represent a long-term accumulation. Since the calcareous shells dissolve in an acidic pH environment in less than two years (cf. Evans 1972) the sample population studied at present very likely includes only specimens born and partly maturing during the several months preceding collection. Having collected the samples in June many of the shells from the previous summer and winter were probably decomposed. Since the samples were collected in early summer there may be some effect on the percentage of juveniles since not all of the species mature at the same time nor at the same rate.

Nearly 65 percent of the identifiable shells were classified as juveniles. Many of the juvenile shells are small *Punctum minutissimum*. If there is a phenological effect on the assemblage this species may be inferred to mature during late June and July although an unknown number of these shells may have died just prior to collection.

Seasonal periodicities are 'nearly universal in communities and often result in almost complete change in community structure during the annual cycle' (Odum 1971: 157). Seasonal changes in abundance of aquatic snails and in the variance of their species abundance distributions were documented for five watercourses in Egypt (Hairston 1964). Critical periods of environmental stress caused the frequency of the most abundant species to decrease drastically; less abundant species underwent a period of increase in many cases at the same time. Strandine (1941) found that fluctuations in a population of *Succinea ovalis* Say coincided with seasonal fluctuations in soil moisture, organic matter and pH.

An alternative conclusion is also possible with respect to phenological variation. Late spring conditions during 1972 may have been deleterious to *Punctum minutissimum*; many of them having succumbed before maturity (cf. Douglas 1963). This can be tested in the future by analyzing additional assemblages under similar conditions and at different times of the year.

The second problem is the taxonomic assignment of *Strobilops labyrinthica*. Identification of species in the genus *Strobilops* are based on shell morphology and, more importantly internal soft part anatomy. *Strobilops labyrinthica*, *S. affinis*, and *S. aenea* are highly variable with respect to shell morphology particularly the number and size of basal folds and lamellae as seen through the base of the shell (Pilsbry 1948).

All of the complete mature specimens from the assemblage thought to be one or the other of these species were measured. None of the specimens was larger than 2.7 mm, which is the minimum diameter of *S. affinis* (Pilsbry 1948). However, none of the

44 specimens was less than 2.5 mm, the maximum size of *S. labyrinthica* (Ibid.)

In addition the shell morphology, other than diameter, suggests that none of the specimens are *S. aenea*; e.g. the lack of an angular periphery characteristic of the latter, which Pilsbry (1948: 862) emphasizes is, 'distinctly but bluntly angular.' The diameter of this species varies from 2.4 to 2.8 mm and thus overlaps the upper limit of *S. labyrinthica* and the lower limit of *S. affinis*.

It thus appears that the specimens of *Strobilops* from this particular assemblage are large examples of *S. labyrinthica* since only one is near the lower limit of *S. affinis* and since all lack the distinguishing morphological characteristics of *S. aenea*.

Analysis of living material of the family Strobilopsidae would provide a more solid basis for specific identification than presently exists on the basis of shell morphology (van der Schalie, personal communication, November 1975).

DESCRIPTIVE STATISTICS

Assessment of variability

The raw frequency data for the gastropod assemblage represented by the first series of 16 samples are summarized in Table 1. Seventeen species of terrestrial gastropods representing thirteen genera and 1,544 identified individuals were recovered. Of these 1,051 were classified as identifiable juveniles (65.1 percent). Juvenile shells of *Punctum minutissimum* contribute in large part to the high percentage of identifiable shells.

There is an apparent high degree of variation in the number of specimens between species; ranging from one specimen of *Gastrocopta corticaria* to 459 specimens of *Punctum minutissimum*. The number of shells recovered from specific samples is also highly variable; ranging from 13 individuals in samples 5 and 6, to 281 individuals in sample 3.

The random manner in which the samples were collected and the size of the assemblage preclude a cursory examination of the raw frequency data in order to assess the major source of variability. The analysis of variance for a randomized complete-block design (Steel and Torrie 1960:134) was used to determine if arithmetic means of species between samples differed significantly, and thus to determine the major source of variation. Due to the apparent lack of a normal distribution of species within or between samples, and to the presence of numerous raw frequencies of less than ten, a transformation of the raw data by the V_{x+1} was used prior to the analysis of variance (Steel and Torrie 1960: 157).

The analysis of variance for the transformed data matrix (Table 2) is summarized in Table 3. The very highly significant ($P > .01$) F ratio for samples (blocks) and species (treatments) indicates, 1) the presence of significant differences of means of spe-

Table 1. Raw frequency of gastropods for each sample. Grid location numbers may be used to locate sample in Figure 1.

Grid Location	1 4R20	2 6R32	3 4SR9	4 9R8	5 16R17	6 18R29	7 19R41	8 21R5	9 28R18	10 30R30	11 31R42	12 33R6	13 40R21	14 42R33	15 43R45	16 7R44	Total
Sample No.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	
<i>Anguispira alternata</i>	2	2	3	4	1		3	6	1			5	3	3	1		34
<i>Euconulus fulvus</i>	-	-	2										1				3
<i>Gastrocopta armifera</i>	-	-						3									3
<i>G. contracta</i>	2												2	1			5
<i>G. corticaria</i>	-	-					1										1
<i>G. pentodon</i>	6	6	29		2	1	2	27	1	4	6	13	22	35	35	20	209
<i>Hawalia minuscula</i>	19	8	29	4	1	10	13	8	15	17	7	16	34	32	24	19	256
<i>Helicodiscus parallelus</i>	16	7	90	1	2	1	4	43	7		2	11	20	35	7	22	268
<i>H. singleyanus</i>	1	6	10		4			32		1				9	8	4	75
<i>Mesodon thyroidus</i>	-	-	1														1
<i>Mesomphix friabilis</i>	-	-						1					5			1	7
<i>Punctum minutissimum</i>	10	21	53	1			4	5		11	4	15	110	108	42	75	459
<i>Pupoides albilabris</i>	-	-											1				1
<i>Retinella indentata</i>	1	3	18	2			1	3				5	11	20	5	8	77
<i>Strobilops labyrinthica</i>	3	9	24	4		1	4	14	1	3	4	13	13	14	7	21	135
<i>Succinea</i> sp.	-	-	3	2			1	1									7
<i>Vallonia perspectiva</i>	-	-			1		1							1			3
TOT. IDENT. IND.	60	62	262	18	11	13	34	143	25	36	23	78	222	258	129	170	1544
Unident. juv.	1	5	7		2		1	2	2			3	7	7	7	3	47
Unident. Fragments	2	2	12	1			1	9	2			8	18		3	8	66
TOT. INDIVID. REC.	63	69	281	19	13	13	36	154	29	36	23	89	247	265	139	181	1657
% Ident. Juv.	63.3	72.6	73.3	66.7	72.7	30.8	67.6	68.5	72.0	55.6	39.1	43.6	65.3	72.1	69.0	76.5	65.1
No. Ident. Juv.	38	45	192	12	8	4	23	98	18	20	9	34	145	186	89	130	1051
No. of Species	9	8	11	7	6	4	10	10	5	5	5	7	11	10	8	8	

Table 2. Raw frequency data of Table 1 transformed by $\sqrt{x+1}$.

Sample No.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
<i>Anguispira alternata</i>	1.73	1.73	2.00	2.24	1.41	1.00	2.00	2.65	1.41	1.00	1.00	2.45	2.00	2.00	1.41	1.00
<i>Euconulus fulvus</i>	1.00	1.00	1.73	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.41	1.00	1.00	1.00
<i>Gastrocopta armifera</i>	1.00	1.00	1.00	1.00	1.00	1.00	1.00	2.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
<i>G. contracta</i>	1.73	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.41	1.00	1.00	1.00	1.73	1.41	1.00	1.00
<i>G. corticaria</i>	1.00	1.00	1.00	1.00	1.00	1.00	1.41	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
<i>G. pentodon</i>	2.65	2.65	5.48	1.00	1.73	1.41	1.73	5.29	1.00	2.24	2.65	3.74	4.80	6.00	6.00	4.58
<i>Hawalia minuscula</i>	4.47	3.00	5.58	2.24	1.41	3.32	3.74	3.00	4.00	4.24	2.83	4.12	5.92	5.74	5.00	4.47
<i>Helicodiscus parallelus</i>	4.12	2.83	9.54	1.41	1.73	1.41	2.24	6.63	2.83	1.00	1.73	3.46	4.58	6.00	2.83	4.80
<i>H. singleyanus</i>	1.41	2.65	3.32	1.00	2.24	1.00	1.00	5.74	1.00	1.41	1.00	1.00	1.00	3.16	3.00	2.24
<i>Mesodon thyroidus</i>	1.00	1.00	1.41	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
<i>Mesomphix friabilis</i>	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.41	1.00	1.00	1.00	1.00	2.45	1.00	1.00	1.41
<i>Punctum minutissimum</i>	3.32	4.69	7.35	1.41	1.00	1.00	2.24	2.45	1.00	3.46	2.24	4.00	10.54	10.44	6.56	8.72
<i>Pupoides albilabris</i>	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.41	1.00	1.00	1.00
<i>Retinella indentata</i>	1.41	2.00	4.36	1.73	1.00	1.00	1.41	2.00	1.00	1.00	1.00	2.45	3.46	4.58	2.45	3.00
<i>Strobilops labyrinthica</i>	2.00	3.16	5.00	2.24	1.00	1.41	2.24	3.87	1.41	2.00	2.24	3.74	3.74	3.87	2.83	4.69
<i>Succinea</i> sp.	1.00	1.00	2.00	1.73	1.00	1.00	1.41	1.41	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
<i>Vallonia perspectiva</i>	1.00	1.00	1.00	1.00	1.41	1.00	1.41	1.00	1.00	1.00	1.00	1.00	1.00	1.41	1.00	1.00

Table 3. Analysis of Variance for data of Table 2.

Source of Variation	df	SS	MS	F
Blocks (samples)	r - 1 = 15	699.986	46.666	-8.006***
Treatments (species)	t - 1 = 16	1451.982	90.992	-15.568***
Error	(r - 1)(t - 1) = 240	-1398.980	-5.829	
Total	rt - 1 = 271			

*** very highly significant P > .05.

Table 4. Descriptive statistics of soil data.

Soil Sample Number	Variables						
	pH	Organic matter %	Avail. P	Acid Sol. P ppm	Exch. K	Exch. Ca me/100g	Exch. Mg
1	6.1	3.6	39.0	54.0	132	10.31	1.37
2	6.2	6.7	45.0	66.7	182*3	13.44	2.08
3	5.6	4.7	43.5	68.3	147	13.75	2.47
4	6.4	5.9	34.1	52.9	219	16.25	2.15
5	5.7	2.3	12.7	24.5	125	5.94	1.24
6	6.0	3.7	20.5	31.5	150	8.75	1.56
7	5.5	4.7	37.1*1	55.5*2	178	7.03	1.37
8	6.6	5.7	34.4	58.3	171	15.63	2.47
9	6.6	3.5	44.0	60.0	418	7.81	1.86
10	6.2	3.9	43.7	63.2	166	11.88	1.37
11	5.5	7.3	55.0	71.2	185	15.63	3.52
12	6.0	5.0	32.8	50.0	122	12.22	2.47
13	6.8	6.7	26.5	43.5	174	20.63	3.45
14	5.2	5.3	43.3	55.0	231	8.13	2.08
15	5.9	6.7	42.5	65.5	150	15.16	2.96
16	7.0	5.3	38.9	65.0	164	16.88	2.67
n	16	16	16	16	16	16	16
\bar{x}	57.30	81.00	593.00	894.90	2914	199.44	35.09
SS	595.61	440.02	23585.50	51473.85	603598	2747.83	84.82
s	6.08	5.06	37.06	55.31	182.13	12.47	2.19
s ²	2603	1.9972	107.1626	168.89	4858.5167	17.4639	5.244
s	5102	1.4132	10.3519	12.9958	69.7031	4.1770	7.242
s _x	1275	3533	2.5880	3.2409	17.4258	1.0444	1.810
C.V.	0839	2782	2793	2358	3827	3352	3302
CL ⁴⁴ L1	5.81	4.31	31.55	48.39	145.00	10.24	1.80
L2	6.35	5.81	42.58	62.23	219.26	14.70	2.58

* The value shown is the x of the remaining 15 values. These values were originally out of agreement with the remaining values.
 *1 Original value = 91.1
 *2 Original value = 136.1
 *3 Original value = 900
 *4 CL (2t, 05s2)

cies between samples; and, 2) that the precision of the sampling design was greatly increased relative to what could have been expected from a completely random design (cf. Steel and Torrie 1960:136).

Data for the seven soil variables analyzed are summarized in Table 4, which also summarizes the descriptive statistics for each soil variable. The variance (s²), standard deviation (s) and standard error of the mean (s_x) differ markedly between variables. The least variable of the parameters is pH (coefficient of variation - C.V. - of 8.39 percent). Considering the apparent variability between samples and among species (Table 3) high positive correlation between a group of gastropod species and pH in this kind of macrohabitat seems unlikely.

The variability of the remaining six parameters approximates two clusters. The first group includes organic matter, available phosphorus and acid soluble phosphorus. The second group includes exchangeable potassium, exchangeable magnesium and exchangeable calcium. The presence of two groups of soil variables with internally similar coefficients of variation suggests the possibility that a complex of species might covary with one or the other of the groups but that no species by itself would covary strongly with only a single soil variable.

Simple correlation of snails and soils

The major purpose of the present study was to determine the relationship of gastropods to the measured soil variables in a quantitative manner. Correlation coefficients, which measure the relationship between two variables on a -1 to +1 scale, satisfy part of this goal. Such correlation coefficients are independent of the scale of measurement; e.g. soil organic matter can be compared with snail species.

One of theseveral options of the computer program used for the factor which follows this discussion computes correlation coefficients between variables. Due to the non-normal distribution the raw frequency data for gastropods was transformed by the $\sqrt{x+1}$. *Gastrocopta armifera*, *G. corticaria*, *Mesodon thyroidus*, and *Pupoides albilabris* were eliminated from the factor analysis due to their rare occurrence. Transformation of the data for these species results in a lack of arithmetical information. Soil data were not transformed since they have a nearly normal distribution. All data were subsequently normalized for the computation of factors. This results in a new data matrix with a mean of zero and a standard deviation of one.

Correlation coefficients between variables are

Variable	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
1 <i>Anguispira alternata</i>	1.000																			
2 <i>Euconulus fulvus</i>	.216	1.000																		
3 <i>Gastrocopta contracta</i>	.125	.167	1.000																	
4 <i>G. pentodon</i>	.299	.399	.105	1.000																
5 <i>Hamaxia minuscula</i>	.087	.502	.511	.666	1.000															
6 <i>Helicodiscus parallelus</i>	.472	.678	.197	.744	.563	1.000														
7 <i>H. singleyenus</i>	.374	.134	-.210	.654	.059	.639	1.000													
8 <i>Mesomphix friabilis</i>	.186	.378	.487	.344	.380	.248	.021	1.000												
9 <i>Punctum minutissimum</i>	.111	.461	.358	.802	.812	.586	.239	.528	1.000											
10 <i>Retinella indentata</i>	.394	.602	.213	.817	.725	.797	.398	.337	.887	1.000										
11 <i>Strobilops labyrinthica</i>	.414	.528	-.025	.808	.583	.809	.498	.365	.763	.855	1.000									
12 <i>Succinea</i> sp.	.490	.576	-.299	.077	-.027	.472	.272	-.110	-.059	.291	.335	1.000								
13 <i>Valtonia perspectiva</i>	.105	-.174	-.011	-.043	-.120	-.053	.046	-.187	.024	.089	-.193	-.036	1.000							
14 pH	-.002	-.036	.190	-.055	-.009	-.009	.001	.568	.061	-.099	.148	-.115	-.598	1.000						
15 % Organic matter	.191	-.094	-.073	.447	.209	.119	.159	.343	.422	.339	.481	.055	-.338	.089	1.000					
16 Available phosphorus	-.118	-.012	-.022	.215	.314	.169	.075	-.268	.190	.147	.280	.066	-.289	-.150	.496	1.000				
17 Acid soluble phosphorus	-.038	.117	-.149	.330	.319	.290	.230	-.163	.254	.224	.456	.205	-.396	.030	.553	.944	1.000			
18 Exchangeable potassium	-.062	-.136	.265	-.285	.042	-.079	-.154	-.061	-.143	-.112	-.225	-.044	-.079	.211	-.058	.327	.216	1.000		
19 Exchangeable calcium	.188	.324	.025	.444	.255	.238	.118	.620	.446	.343	.582	.142	-.656	.530	.755	.207	.379	-.216	1.000	
20 Exchangeable magnesium	.087	.325	-.046	.562	.330	.293	.125	.522	.502	.428	.551	.006	-.432	.198	.826	.340	.411	-.029	.820	1.000

Table 5. Covariances between variables; includes gastropod and soil variables. Correlation coefficients of snail species are based on all frequencies of each variable transformed by $\sqrt{N+1}$. Coefficients for soil variables are based on raw data of all soil parameters.

summarized in Table 5. The coefficients are significant at the five percent level if they equal or exceed .497 (onn-2 degrees of freedom; Freese 1967: 87). A coefficient equal to or exceeding .623 is significant at the one percent level.

Examination of the correlation coefficients permits the delineation of two groups of gastropod species. The members of each group form a complex of species with relatively similar correlation coefficients; those species which significantly correlate at the five percent level and those which show significant correlation at the one percent level. The correlation coefficients are based on comparisons between two variables at one time and depend on the direction (increase or decrease) of frequencies relative to each other. Quantitative definition of groups of correlated species is difficult on this basis since only two variables are being compared.

Very few significant correlations were obtained between snail species and soil variables. *Valtonia perspectiva* is negatively correlated with soil pH. A negative correlation was not anticipated since *V. perspectiva* is most common in more open calcareous environments (Grimm 1959:22) and would be expected, in this case, to increase as pH increased. The very low frequency of *V. perspectiva* precludes a definite conclusion about the relationship of this species to soil parameters. It should also be noted that this species is significantly negatively correlated with calcium. The calcium values of some samples are very low although available phosphorus levels are quite high (Dr. Theodore Peck, personal communication).

The ratio of available phosphorus to acid soluble phosphorus levels is normally low in acid soils

due to the insolubility of calcium phosphates in acid environments (Thompson and Troeh 1973: 269). A correlation coefficient of .944 between these two variables was obtained in this analysis. Most of the available phosphorus in acid soils is complexed with aluminum. The low frequency of *Valtonia perspectiva* in this situation is likely due to the lack of calcium which would be in the exchangeable form in calcareous environments in open areas where this species is often abundant.

Calcium and magnesium behave similarly in the soil system (Thompson and Troeh 1973: 316); a correlation of .820 was obtained in the present study. It may be expected then that gastropod species should correlate with magnesium and calcium levels in a similar manner. For example in addition to the negative correlation between *Valtonia perspectiva* and calcium this species is also negatively correlated with magnesium, although the correlation coefficient is not significant.

Mesomphix friabilis on the other hand has a significant positive correlation with pH, calcium and magnesium (.568, .620, and .522 respectively). Unfortunately this species occurred in such low frequencies that additional testing is necessary before its relationship to pH can be adequately assessed. One, five and one specimen of *M. friabilis* were recovered from samples 8, 13 and 16 respectively. The pH values for these samples are 6.6, 6.8 and 7.0 respectively. It can be tentatively concluded on this basis that *M. friabilis* is a sensitive indicator of pH ranging from 6.6 to 7.0 in addition to the presence of moist conditions (F.C. Baker 1939: 67). The shell thickness of this species decreases in lime deficient areas (Pilsbry 1946: 330). The soils of Mound 2, with low values of calcium and acidic pH indicate the lack of free

carbonates in solution. The high correlation between this species and exchangeable calcium (.620) and magnesium (.522) suggests that it is existing in a less than optimal environment with a low calcium supply. These data clearly support the conclusions of F.C. Baker and Pilsbry.

Other species have high correlations with magnesium; *Gastrocopta pentodon*, *Punctum minutissimum*, and *Strobilops labyrinthica*. Although correlation coefficients between these species and calcium are positive the only apparently significant correlation is between *Strobilops labyrinthica* and calcium (.582). These three species, in addition to *Mesomphix friabilis*, form a group which is closely related to the supply of calcium and magnesium.

No significant correlations were obtained between the 13 snail species and the soil variables of organic matter, available phosphorus, acid soluble phosphorus and exchangeable magnesium.

A few species are correlated with increases in organic matter but not significantly; *Gastrocopta pentodon*, *Punctum minutissimum*, and *Strobilops labyrinthica*. The correlation between the latter species and acid soluble phosphorus (.456) although not significant is positive.

With the exceptions of *Mesomphix friabilis* and *Vallonia perspectiva* there is generally a nonlinear or slightly negative correlation between snail species and pH.

The use of correlation coefficients as a basis for studying the relationship of gastropods and soil variables substantiates some conclusions reached by previous investigators. The use of this bivariate statistical approach is perhaps most appropriate when dealing with species which may inhabit such macrohabitats as were studied in this case but which are most common in other environments. The data of the present analysis indicate that *Vallonia perspectiva* and *Mesomphix friabilis*, which are most common in other situations, are not common in this particular habitat because their environmental tolerances have been very nearly exceeded. The results give some insight into the environmental requirements of these species and a more detailed understanding of why they are common in their optimal habitats.

Other species, such as *Hawaitia minuscula*, *Helicodiscus parallelus* and *H. singleyanus*, occur in a very wide range of habitats and are common in the assemblage presently under study. In the present analysis these species show a generally positive correlation with all of the measured soil parameters except exchangeable potassium with which they show a negative correlation.

The soil variables analyzed in the present analysis are the same as those examined by Burch (1955) with the addition of acid soluble phosphorus data. For purposes of comparison the values of phosphorus and potassium are expressed in percentages by multiplying parts per million by 10^{-4} . Calcium and magnesium values are expressed in milliequivalents

per 100 grams. These may be converted to parts per million by multiplying the calcium values by 200 and the magnesium values by 120, and expressed in percentages by multiplying the parts per million value by 10^{-4} . It is not necessary to relate all variables to one standard since the variance ratios between variables will remain nearly the same irrespective of the standard.

In contrast to Burch's data exchangeable potassium values are usually higher, and, like the organic matter and exchangeable calcium values, the lower end of the ranges overlaps the upper range of Burch's values. Magnesium values are higher and available phosphorus values are lower than those obtained by Burch. Soil pH values are comparable between the two data sets.

One source of variation in the two data sets is the analytical precision of the various soil testing techniques. A second possible reason for the differences between several of the soil variables is the variation in climate between the two regions, in addition to other soil factors specifically parent material and time. A third important source of variation is the difference in sample size. Burch utilized a regional approach obtaining soil data from 41 stations, encompassing two physiographic regions, the Coastal Plain and Piedmont Plateau. All of these differences preclude a direct comparison between the two data sets, although it is tempting to use the values obtained in this study to suggest the possible relationship between snails and some soil parameters, for example, at higher levels of organic matter content.

Burch plotted soil parameters against the number of species and secondly against the number of specimens. In doing so he used an evenly spaced scale but each increment of the scale did not in all cases represent an equal increase in values of the same variable along the scale. The result is a graphic representation of relationships between the number of species and soil variables and between the number of snail specimens and soil data.

The present study utilizes a much different approach for graphical representation of the data. Descriptive statistics were obtained for the raw soil data (Table 4). A frequency distribution was constructed for each variable using the respective mean as the midpoint and the appropriate standard deviation as the increment. These data and the frequency distributions showed all soil variables, except exchangeable potassium, to have a near normal distribution. To compare these data with those of Burch an even increment was used. To obtain a similar scaling effect for all parameters the range of each was divided into six equal parts and plotted against the number of species, then against the number of specimens.

Inspection of Figures 2A and 2B shows some relationships between the number of species and soils and between the number of specimens and soil data, although in both sets of graphs there is a great deal of variation. Rather than illustrating causal affinities (the 'relationships' of Burch) between




Figure 2A. Even increment plot of soil data against the number of small specimens.




Figure 2B. Even increment plot of soil data against the number of small specimens.




Figure 3A. Plot of soil data against number of species using standard deviation as the increment.




Figure 3B. Plot of soil data against number of specimens using standard deviation as the increment.

Table 6. Percent of total variance explained by respective eigenvector.

Eigenvalues ¹	Fraction of Total Variance	
	Individual ²	Cumulative
7.12823	35.64	35.64
2.94875	14.74	50.38
2.49691	12.48	62.87
1.97363	9.87	72.74

1 Eigenvalue = sum of the column of squared loadings for each factor. Measures the amount of variance accounted for by a factor.

2 $\frac{\text{Eigenvalue}}{\text{number of variables}} \times 100 =$ Percent of total variance explained by the eigenvector.

snail species and soil variables. graphs such as these represent evidence of the optimal habitat of species within a range of given soil parameters; i.e. how diversity and frequency relate to a range in soil characteristics. The graphs consider all species present rather than ecological dominants or covariation between numbers of species or numbers of specimens, and soil parameters. This occurs due to the near-normal distribution of the values for nearly all of the soil variables. Most soil values are within one standard deviation of the mean value. Thus most of the snail samples also lie within these limits; i. e. the soil samples and the snail samples are the same points in space. This unavoidably increases the area and thus the number of species since the latter tends to increase in any population as the area of the sample increases (Odum 1971: 143-154). The use of an increment less than one standard deviation would even out the sharp inflections in the distribution curves. This however defeats the purpose of attempting to delineate patterns of relationship between gastropods and soils on a quantitative basis.

A pattern similar to even increment scaling is obtained when the soil data are plotted against the number of species and the number of specimens using the standard deviation of each variable as the increment (Figures 3A and 3B).

The use of the above analytical techniques facilitates interpretations about the assemblage as a whole but permits very few inferences about autecology and the significance of particular species in a particular habitat.

In conclusion, the optimal conditions in this habitat utilizing Figures 3A and 3B, considering the complex of soil variables measured, for nine to thirteen of the species represented, are characterized by exchangeable calcium levels of 8.29 to about 20.83 milliequivalents per 100 grams of magnesium. Acid soluble phosphorus levels would be from between 42.31 and 44.31 parts per million to less than 81.31 parts per million. Soil pH would

range from 5.57 to about 7.0, and organic matter from 3.65 to 6.47 percent. This 'habitat' would predictably contain from 350 to around 700 individuals considering a sample size of about nine square feet.

Multivariate analysis

The univariate and bivariate statistics described previously are not sufficient to quantify a relationship between several variables which may be related in a multivariate and complex manner.

The principal tool used to delineate the quantitative covariation between snail species and soil variables discussed in this study was multivariate factor analysis. The computer program for the analysis was written in FORTRAN by Mr. Peter Guetter of the Department of Meteorology of the University of Wisconsin. The program was initially applied to climatic data but is applicable to a wide range of data sets. Included in the program is the generation of a number of factors here to be referred to as eigenvectors. The factors were generated from the data in Table 2; snail frequencies transformed by the $\sqrt{x+1}$, and soil parameters. Each eigenvector 'summarizes' the mathematical relationships between a number of variables; i.e. the covariance of interrelated variables. As such the analysis implies no statistical significance (Rummel 1967). The intent of factor analysis is to 'express covariation in terms of k underlying factors that explain a large part of the variance and covariance of the original variables. The number of factors is much less than that of the number of variables in the study' (Sokal and Rohlf 1969: 542).

Eigenvalues measure the amount of mathematical variance accounted for by an eigenvector. The eigenvalues for the present analysis and the variance they account for, expressed as a percent, are summarized in Table 6. Fifteen eigenvectors were computed; those which explain less than ten percent of the total amount of variability (100 percent) are considered insignificant.

Table 7. Eigenvectors with factor loadings for each variable.

Variable	Eigenvector			
	1	2	3	4
1 <i>Anguispira alternata</i>	-.140406	-.222126	.005389	-.270861
2 <i>Euconulus fulvus</i>	-.232059	-.163953	-.066870	-.077894
3 <i>Gastrocopta contracta</i>	-.074472	.001533	-.400785	.352647
4 <i>Gastrocopta pentodon</i>	-.324577	-.131144	.022044	.053415
5 <i>Hawaiiia minuscula</i>	-.266096	-.058836	-.134704	.383516
6 <i>Helicodiscus parallelus</i>	-.294690	-.264080	.051647	-.014480
7 <i>Helicodiscus singleyanus</i>	-.170687	-.209209	.196635	-.196011
8 <i>Mesomphix friabilis</i>	-.195757	.122171	-.461717	-.113591
9 <i>Punctum minutissimum</i>	-.311730	-.068208	-.158576	.235900
10 <i>Retinella indentata</i>	-.321627	-.224146	-.032492	.113066
11 <i>Strobilops labyrinthica</i>	-.344220	-.076658	.069540	-.075626
12 <i>Succinea</i> sp.	-.105983	-.201880	.253882	-.327742
13 <i>Vallonia perspectiva</i>	.107645	-.406303	-.012034	.182406
14 pH	-.061170	.316205	-.286369	-.223062
15 % Organic matter	-.230443	.297354	.140053	-.069531
16 Available phosphorus	-.129625	.235334	.420607	.335622
17 Acid soluble phosphorus	-.182229	.239256	.424116	.180917
18 Exchangeable potassium	.049858	.148663	.041001	.315285
19 Exchangeable calcium	-.260216	.326115	-.077150	-.275137
20 Exchangeable magnesium	-.267310	.280348	-.006743	-.069492

The first eigenvector explains 35.6 percent of the total variance in the data set of 20 variables. Each successive eigenvector explains a lesser portion of the remaining variance. The factor loadings for each variable (Table 7) indicate a general pattern of negative covariance between variables. The factor loadings being nearly all negative force more of a subjective approach toward emphasizing the variables which are important in the first eigenvector. *Gastrocopta pentodon*, *Punctum minutissimum*, *Retinella indentata*, and *Strobilops labyrinthica* are the variables which have the greatest similarity of covariance as expressed by factor loadings. Much of the variance is explained by the first eigenvector (Table 8: 75, 69.3, 72.7, and 84.5 percent respectively). The factor scores and the frequent occurrence of these species in most of the 16 samples provide the basis for the tentative conclusion that these species characterize the forest habitat.

Factor loadings for samples (Table 9) show that samples 3 and 13 are quite similar with respect to the information they contribute to the first eigenvector. These samples in addition to samples 8, 14, and 16 covary in a negative fashion with a similar amplitude, opposite to samples 5, 6, 7, and 9. These results indicate substantially different patterns of variation between the two groups of samples.

A second group of variables has a high degree of covariance in the first eigenvector. These variables are: *Hawaiiia minuscula*, *Helicodiscus parallelus*, calcium, and magnesium. Although these have no significant linear correlations with each other (Table 5) they are summarized in the first eigenvector (Table 8).

The two groups of covarying species evident in the first eigenvector contain species which occur

Table 8. Percent of explained variance for each variable, first four eigenvectors.

Variable	Eigenvector				Total
	1	2	3	4	
1 Anugispira alternata	14.05	14.55	.00	14.48	43.08
2 Euconulus fulvus	38.39	7.93	1.12	1.20	48.64
3 Gastrocopta contracta	3.95	.00	40.11	24.54	68.60
4 Gastrocopta pentodon	75.10	5.07	.12	.56	80.85
5 Hawalia minuscula	50.47	1.02	4.53	29.03	85.05
6 Helicodiscus parallelus	61.90	20.56	.67	.04	83.17
7 Helicodiscus singleyanus	20.77	12.91	9.65	7.58	50.91
8 Mesomphix friabilis	27.32	4.40	53.23	2.55	87.50
9 Punctum minutissimum	69.27	1.37	6.28	10.98	87.90
10 Retinella indentata	73.74	14.81	.26	2.52	91.33
11 Strobilops labyrinthica	84.46	1.73	1.21	1.13	88.53
12 Succinea sp.	8.01	12.02	16.09	21.20	57.32
13 Vallonia perspectiva	8.26	48.68	.04	6.57	63.55
14 pH	2.67	29.48	20.48	9.82	62.45
15 % Organic matter	37.85	26.07	4.90	.95	69.77
16 Available phosphorus	11.98	16.33	44.17	22.23	94.71
17 Acid soluble phosphorus	23.67	16.88	44.91	6.46	91.92
18 Exchangeable potassium	1.77	6.52	.42	19.62	28.33
19 Exchangeable calcium	48.27	31.36	1.49	14.94	96.10
20 Exchangeable magnesium	50.93	23.18	.01	.95	75.07

in nearly all of the samples and which have the highest frequencies. These species are concluded to be characteristic of the habitat of the sample area. The lack of significant correlations (Table 5) between these species and soil variables, and the lack of evident covariance based on factor scores, suggests that they are responding predominantly to other ecological parameters. At least fifty percent of the variance of these species is explained by the first eigenvector.

Much of the remaining variance of *Helicodiscus parallelus* and *Retinella indentata* is explained by the second eigenvector (Table 8). With much of their variance already accounted for, these species and *Anguispira alternata* and *Helicodiscus singley-*

anus show a general pattern of covariance of similar amplitude as six of the seven soil variables (excluding potassium). The covariance of the snail species, however, is in a direction opposite that of the soil variables. Samples 1, 4, 9, 10, 11, 13, and 16 contribute most of the variance accounted for by the second eigenvector (Table 9). These samples also suggest a pattern of distribution of the variance between sample points that is opposite to the pattern indicated by samples 3, 5, 6, 7, and 14.

Following the computation of the second eigenvector 50.4 percent of the total variance of the assemblage has been accounted for (Table 6). In the third eigenvector, the last one considered to be of

Table 9. Eigenvectors with factor loadings for each observation (sample).

Sample	Eigenvector			
	1	2	3	4
1	1.208321	-.336089	-1.056477	1.189470
2	-.126662	1.225691	1.281039	-.228042
3	-4.692862	-2.754288	1.911508	-.595574
4	1.293829	1.070858	.639006	-2.156628
5	4.813039	-2.822209	-1.419051	-1.193376
6	3.621002	-.239646	-1.288072	-.644324
7	2.281698	-1.696686	.931875	.410141
8	-2.053272	-.516163	.747936	-2.749899
9	2.520461	1.487161	-.207997	2.112110
10	1.920869	1.073838	.487117	.765144
11	.363201	3.011929	2.092990	.548227
12	-.391342	-.148436	-.191281	-.858923
13	-4.349376	1.190759	-4.685825	-.120355
14	-2.302820	-2.963130	.299530	3.031444
15	-1.735203	.998141	.997426	.497122
16	-2.370887	1.418273	-.539726	-.006536

pertinent value in this analysis, *Gastrocopta contracta* and *Mesodon friabilis* have factor scores of similar amplitude but which covary negatively with available and acid soluble phosphorus (Table 5). The remaining species tend to covary negatively or to have little relationship to soil variables.

The results of the eigenvector analysis, in addition to the results of the simple correlations (Table 5), permit the conclusion that the soil parameters measured in this study are of secondary or tertiary importance with respect to the distribution of the species included in the factor analysis.

Qualitative habitat differentiation

During field study the sample area appeared to contain two habitats which could be distinguished on the basis of apparent cover; a sparse grass-and-moss association on top of the mound and a humus layer increasing in thickness on the sides and toe-slope of the mound. It was hypothesized that the analytical method used to study covariance of species could provide data justifying a qualitative distinction between the two habitats.

Inspection of the transformed data matrix (Table 2) and Figure 1 permit the conclusion that the two

habitats do contain markedly different data sets with respect to snail frequencies and soil parameters. On close inspection a third habitat is distinguishable. The east slope of the mound differs from the west slope; the former has a higher frequency of *Punctum minutissimum* and *Gastrocopta pentodon*, a generally higher frequency of other species (excluding species not included in the factor analysis), and a generally greater organic matter content per unit area. The samples included in arriving at this conclusion are 3, 13, and 14.

The top of the mound (samples 5, 6, 9, and 10) is separable from the other microhabitats due to generally lower snail frequencies, particularly of *Gastrocopta pentodon* and *Punctum minutissimum*, and lower organic matter content.

Due to the transformation of the raw frequency matrix a conclusion regarding differences in species diversity between habitats must be based on Table I. Again, the samples from the top of the mound are conspicuous because of the fewer species per unit area than samples from the west side of the mound. Samples from the east side of the mound are also conspicuous because the species diversity of these is greater than in any of the remaining samples (Table I).

The habitat differentiation is also supported by the eigenvector analysis. *Hawaiiia minuscula*, *Helicodiscus singleyanus*, and *H. parallelus* tend to predominate among the species present in the samples from the top of the mound (Table 7 and Figure 1). *Hawaiiia minuscula* and *Helicodiscus parallelus*, in addition to magnesium and calcium, had a high degree of covariation in the first eigenvector. This is due in part to the pattern of variation of these variables in samples from the top of the mound, even though the same species occur in other samples in higher frequencies. Other species found to covary in the first eigenvector included *Gastrocopta pentodon*, *Punctum minutissimum*, *Retinella indentata*, and *Strobilops labyrinthica*. These species occur in all of the samples but attain their highest frequencies in samples 3, 13, and 14; i. e. on the east side of the mound.

The previous qualitative distinctions give further indication that microhabitat differentiation within a macrohabitat is possible and is quantifiable with this form of eigenvector analysis, with the advantage of quantitative definition of relationships between many variables simultaneously.

SUMMARY AND CONCLUSIONS

The species which tended to covary throughout the eigenvector analysis were those which occurred in all or nearly all of the 16 samples and which had the highest raw and normalized frequencies. This group, characteristic of the sample area, includes *Gastrocopta pentodon*, *Punctum minutissimum*, *Retinella indentata*, and *Strobilops labyrinthica*. Additional species form a group of secondary importance but the members still covary; *Hawaiiia minuscula* and *Helicodiscus parallelus*. *Anguispira al-*

ternata and *Helicodiscus singleyanus* form a group also of secondary importance.

Species included in the primary group have been shown by numerous investigators to be characteristic of forested habitats (Burch 1956; Douglas 1963; Atkins 1966; Elwell and Ulmer 1971). Nearly all of the remaining species in the assemblage are common in other forested habitats, with the exception of *Vallonia perspectiva* and *Gastrocopta armifera*. All of the species of this assemblage could be expected to occur in a range of habitats; i. e. many species may occur in areas more open than the Mueller-Ringhausen forest, particularly *Hawaiiia minuscula*, *Helicodiscus parallelus*, *Vallonia perspectiva*, and *Gastrocopta armifera*.

Previous authors (Burch 1955, 1956) have concluded that potassium, phosphorus and pH have an indirect effect on the distribution of land snails. Burch's study considered the relationships of soil variables and snails on a regional level. At the local scale of the present analysis two species were found to have significant simple correlation with soil pH; *Mesomphix friabilis* and *Vallonia perspectiva*, the latter having a negative correlation. *Mesomphix friabilis* was also significantly correlated with calcium and magnesium.

The latter species, in addition to *Gastrocopta pentodon*, *Punctum minutissimum*, and *Strobilops labyrinthica*, were highly correlated with magnesium. Using multivariate techniques, however, i. e. examining all the variables simultaneously, there was no definite pattern of covariance between any of the above species, which were the most frequently occurring ones, and soil variables. When examined bivariately these species appear to be responding to the pattern of variation in principally magnesium, but also to calcium.

There is an absence of significant bivariate correlations between other species and organic matter, available phosphorus and exchangeable magnesium. The eigenvector analysis showed all of the soil variables to be of secondary importance with respect to their effect on the distribution of snails. The major covarying variables are complexes of species rather than species and soil variables.

These results also disagree with Burch's conclusion regarding the effect of organic matter on the distribution of snails. In the present study organic matter is concluded to be of minor importance in terms of its covariance with a complex of snail species characteristic of the local area.

During the preceding analysis an attempt was made to assess qualitatively the relationships between gastropods and several soil variables. A quantitative approach was then used to delineate statistically significant relationships in several cases where two variables were examined together. A factor analysis was utilized to summarize the multivariate relationships between 20 variables, 7 of which were soil parameters.

The humid temperate weathering regime of the Illinois Valley and surrounding regions is the pri-

mary factor affecting the development and maintenance of the expansive deciduous forests of the Prairie Peninsula. It is in the deciduous forests, where soils are acidic and on acidic soils of transition zones between forests and prairie, where the quantitative ecology of land snails adapting to soil, cover, and moisture conditions will be most substantially understood. This is due primarily to the lack of significant effects brought about by the preservation of shells, as happens in alkaline environments. In addition, refined sampling techniques can be expected to result in qualitative measurement of intra- and inter-seasonal variation in the molluscan assemblages, and thus in quantitatively based analyses of changes in gastropod communities.

The quantitative definition of habitat groups, similarity of covariance of a complex of species, has distinct advantages over subjective assessment of species lists and the comparison of only a few species at a time. Continued use of multivariate techniques and refined data collecting techniques will contribute to a more complete understanding of the relationship between land snails and their physical environment. Such improvements will facilitate more detailed interpretations of past molluscan assemblages, and thus more accurate reconstructions of past environments based on terrestrial mollusks.

LITERATURE CITED

- ATKINS, C.G. (1966) Factors affecting the structure and distribution of terrestrial Pulmonata. -- Proc. Iowa Acad. Sci. 73: 408-416
- BAKER, F. C. (1939) Fieldbook of Illinois land snails. -- Manual 2, Ill. Nat. Hist. Survey, Urbana
- BLACK, C. A. (1965) Methods of soil analysis. Part 2: Chemical and microbiological properties. -- Amer. Soc. Agron., Madison
- BOYCOTT, A. E. (1929) The oecology of British land Mollusca, with special reference to those of ill-defined habitat. -- Proc. Malac. Soc. London 18 (5): 213-224
- BRAY, R. H. & KURTZ, L. T. (1945) Determination of total, organic, and available forms of phosphorus in soils. -- Soil Sci. 59: 39-45
- BURCH, J. B. (1955) Some ecological factors of the soil affecting the distribution and abundance of the land snails of eastern Virginia. -- Naut. 69 (2): 62-69
- (1956) Distribution of land snails in plant associations in eastern Virginia. -- Naut. 70 (2): 60-64
- (1962) How to know the eastern land snails. -- Wm. C. Brown Co.
- CURTIS, J.T. (1959) The vegetation of Wisconsin. -- University of Wisconsin Press, Madison
- DOUGLAS, C. I. (1963) Population analyses, variation and behavior of *Anguispira alternata alternata*. -- Trans. Kansas Acad. Sci. 66 (2): 186-194
- ELWELL, A. & ULMER, M.J. (1971) Notes on the biology of *Anguispira alternata* (Stylommatophora: Endodontidae). -- Malacologia 11 (1): 199-215
- EVANS, J. G. (1972) Land snails in Archaeology. -- Seminar Press
- FREESE, F. (1967) Elementary statistical methods for foresters. -- Handbook 317, U.S.D.A., Washington
- GETZ, L. L. (1974) Species diversity of terrestrial snails in the Great Smoky Mountains. -- Naut. 88 (1): 6-9
- GRIMM, W. (1959) Land snails from Maryland and Virginia. -- Naut. 73 (1): 21-22
- HAGGETT, P. (1965) Locational analysis in human geography. -- Edward Arnold Publ. Co.
- JAEHNIG, M.E. W. (1971) The study of gastropods: methodology. -- Plains Anthropologist 16 (54) Part 1: 289-297
- ODUM, E. P. (1971) Fundamentals of ecology. -- Third edition; W. B. Saunders Co.
- OUGHTON, J. (1948) A zoogeographical study of the land snails of Ontario. -- Univ. Toronto Studies, Biol. series 57
- PILSBRY, H. A. (1946) Land Mollusca of North America (North of Mexico). -- Monograph 3, v. 2 (1), Acad. Nat. Sci. Philadelphia
- (1948) Land Mollusca of North America (North of Mexico. -- Ibid., v. 2 (2)
- RIGGLE, R. S. (1975) Quantitative definition of prehistoric and contemporary molluscan habitat groups in the Lower Illinois River Valley. -- Unpubl. M.A. thesis (Anthropology), Univ. Wisconsin, Madison
- RUMMEL, R. J. (1967) Understanding factor analysis. -- J. Conflict Resolution 4: 444-480
- SHIMEK, B. (1930) Land snails as indicators of ecological conditions. -- Ecology 14: 673-686
- SOKAL, R. R. & ROHLF, F. J. (1969) Biometry. -- W. H. Freeman & Co.
- STEEL, R. G. D. & TORRIE, J.H. (1960) Principles and procedures of statistics. -- McGraw-Hill Book Co., Inc.
- STRANDINE, E. J. (1941) Quantitative study of a snail population. -- Ecology 22 (1): 86-91
- THOMPSON, L. M. & TROEH, F. R. (1973) Soils and soil fertility. -- Third ed., McGraw-Hill Book Co., Inc.
- VAN CLEAVE, M. J. (1951) Snails in their relations to the soil. -- The Biologist 34: 3-9

POSTSCRIPT ON MUSSEL VS. MUSCLE
IN RELATION TO BIVALVES

RALPH W. DEXTER

Kent State University, Kent, Ohio

Confusion in the use of the common word 'mussel' or 'muscle' in reference to bivalve mollusks has been an issue of long standing. The problem has involved usage in publications and on maps, and especially with reference to what is officially recognized as the 'Muscle Shoals' in Tennessee. Writers especially concerned over this matter have been Ortman (1924), Matthes (1925), Dexter (1961, 1967) and Isom (1971). Since my last report I have encountered a number of additional cases of sufficient interest to warrant publication.

Daniel Drake (1815) in his historically important account of Cincinnati and its environs consistently refers to 'muscle shells' in reference to the clams of that area. According to H. E. Wheeler (1935), Timothy Abbott Conrad wrote to Judge Charles Tait on 10 April 1834, 'I have my boxes safely deposited in my bedroom, the floor of which is carpeted with muscles and fossils not greatly to the edification of my mother and sisters.' Col. Charles Whittlesey (1869) refers to the 'muscle Shoals' in the paper he read at the meetings of the American Association for the Advancement of Science.

I have long been concerned about the spelling of Mussel Point at Cape Ann, Mass. As noted earlier the oldest maps and atlases used the obsolete spelling 'muscle Point,' but on the modern maps it is given as Mussel Point. One exception is the U. S. Coast and Geodetic Survey Map 243, East Coast Mass., Ipswich Bay to Gloucester Harbor, published in 1958, which reverted to the obsolete spelling. An inconsistency was noted on the U. S. Geological Survey 'Atlas of the City of Gloucester and Town of Rockport, Mass.' published in 1884 in which the obsolete spelling appears in the Atlas but the correct spelling 'Mussel Point' appears on the map.

In Jordan's report on the California earthquake (1907) he makes reference to Mussel Rock on the shore of San Francisco. In modern maps of the Indian Ocean, Mussel Island is designated in the Gulf of Manaar of the Indian Ocean. On the other hand, the Reverend Henry W. Winkley of Saco, Maine, wrote to Dr. Victor Sterki 19 Nov. 1891, 'the last item in the *Nautilus* speaks of the fish *Rhodius amarus* depositing eggs in the mantle of freshwater muscles, etc.' Dr. Victor Sterki, of New Philadelphia, Ohio, wrote to Dr. Henry A. Pilsbry, Curator of Mollusks at the Academy of Natural Sciences of Philadelphia in Pennsylvania and editor and publisher of the *Nautilus*, 20 March 1899 as follows, 'If the word

mussel was changed to muscle it was a printer's blunder, which I overlooked when reading proof. Your use of the term (mussel) is of course perfectly correct.' (Both letters at the Carnegie Museum, Pittsburgh, Pa.) Possibly changes made by typesetters and printers may explain some of the confusion in the use of these words.

In Sterkiana No. 49 (March, 1973) Isom (p. 18-20) criticized Stansbery for continuing to use the name 'Mussel Shoals' in his report on 'Rare and Endangered Freshwater Mollusks in Eastern United States.' Officially the locality is known as 'Muscle Shoals,' but some malacologists continue to use the more appropriate spelling. Although van der Schalie in Sterkiana No. 52 (1973) hedges by using the official name, he puts quotation marks around it (i.e. 'Muscle Shoals').

LITERATURE CITED

DEXTER, R. W. (1961) Mussel Shoals vs. Muscle Shoals. -- Sterkiana No. 4: 29-31

---- (1967) Addenda on Mussel vs. Muscle. -- Sterkiana No. 27: 1-2

DRAKE, Daniel (1815) Natural and statistical view or picture of Cincinnati and the Miami Country. -- 251 p.

ISOM, B. G. (1971) A biologist's look at the history of Muscle Shoals (Mussel Shoals). -- Malacological Review 4: 203-206

JORDAN, D. S. (ed.) (1907) The California Earthquake of 1906. -- 371 p.

MATTHES, G. H. (1925) Muscle Shoals vs. Mussel Shoals. -- Science 61: 209

ORTMANN, A. E. (1924) Mussel Shoals. -- Science 60: 565-566

van der SCHALIE, H. (1973) The mollusks of the Duck River drainage in central Tennessee. -- Sterkiana No. 52: 45-55

WHEELER, H. E. (1935) Timothy Abbott Conrad. -- Bull. Amer. Paleont. 23: 54

WHITTLESEY, C. (1869) On the evidence of the antiquity of Man in the United States. -- Proc. Amer. Assoc. Adv. Sci. 17: 268-288

Accepted for publication December 20, 1975

MICRO-DISTRIBUTION OF LAND SNAILS IN AN ARTIFICIAL TALUS SLOPE

RAYMOND W. NECK

Texas Parks & Wildlife Department
John H. Reagan Building
Austin, Texas 78701

An examination of an artificial talus slope in Alpine, Brewster Co., Texas, on 16 December 1974 revealed a heterogeneous environment in relation to suitability of microhabitats for land snails. This man-made talus slope formed an edge for a fill area which had been raised to a level plane from its natural slope condition. Two types of rocks were used in the fill material (apparently at random as no intentional layering could be detected). Rocks included a light-colored aphanitic igneous rock and a hard fossiliferous limestone.

Four species of snails were observed at this locality: Achatinidae: *Rumina decollata* L. (54 living snails and numerous dead snail shells); Pupillidae: *Gastrocopta procera procera* (Gould) (38 living snails) and *Pupoides albilabris* (C. B. Adams) (1 living snail); and Succineidae: *Succinea* sp. (*grosvenori* Lea?) (4 empty shells). *R. decollata* and *P. albilabris* have previously been reported from Brewster County (Cheatum *et al.* 1972). This collection is a new county record for *G. p. procera*, although it is known from three of the four adjoining counties (Cheatum *et al.* 1972; Leonard & Frye 1962). The *Succinea* cannot be positively identified as no live material could be found. Shells collected at this site are similar to those of *S. grosvenori*. *S. grosvenori* has not been reported from Brewster Co., although it is known from adjoining Jeff Davis Co. (Cheatum *et al.* 1972).

Distribution of the snails under the rocks of the talus slope is summarized in Table 1. The data reveal a decided preference for the limestone rock as a source of resting spots. An analysis of the data by X-square test reveals the difference to be significant at greater than the 0.001 level. The application of the test in this case is marginally proper as the expected value for one of the categories is less than five (but only slightly at 4.7). Utilization of Yates' correction negates most of this problem. At any rate, the data are obviously significant by visual examination of Table 1. No snails were found under the igneous rocks.

Shelled mollusks are greatly affected by soil pH and calcium ion concentration (see reviews by Lozek 1962; Newell 1967). In this case, however, the soil is homogeneous; some factor associated with the rocks involved affected snail distribution. If one assumes that both types of rocks provide

sufficient protection from desiccation and overheating from excessive solar radiation, some chemical factor(s) is (are) the determinant(s) of the distribution of these snails.

One would expect finding a surplus of snails in association with the limestone rocks. The absence of snails under the igneous rocks indicates the likelihood of the existence of a repellent character which apparently prevents these snails from using those rocks as resting sites.

REFERENCES

- CHEATUM, E.P., FULLINGTON, R., & PRAIT, L. (1972) Molluscan records from West Texas. -- *Sterkiana* 46: 6-10
- LEONARD, A. B. & FRYE, J. C. (1962) Pleistocene molluscan faunas and physiographic history of Pecos Valley in Texas. -- *Bur. Econ. Geol., U. Texas, Rept. Invest.* 45, 42 p.
- LOZEK, V. (1962) Soil conditions and their influence on terrestrial Gasteropoda in central Europe. -- IN: *Progress in soil zoology*, P.W. Murphy, Butterworth, p. 334-342
- NEWELL, P.F. (1967) Mollusca. -- IN: *Soil biology*, A. Burges & F. Raw, Academic Press, p. 413-433.

TABLE 1. DISTRIBUTION OF LAND SNAILS UNDER ROCKS

SNAILS	ROCK TYPES		TOTAL
	igneous	limestone	
present	0*	11	11
absent	19	3	22
TOTAL	19	14	33

* Numbers refer to number of rocks in each category.

$X^2 = 18.99$ d.f. = 1 $p < 0.001$

Accepted for publication January, 1976

TESTACELLA HALIOTIDEA (DRAPARNAUD)

FREE LIVING IN PENNSYLVANIA

BRANLEY A. BRANSON

*Eastern Kentucky University
Richmond, Kentucky 40475*

On 9 October 1975, Dr. C. J. McCoy, Jr., Carnegie Institute, Pittsburgh, Pennsylvania, collected one adult specimen of this slug from the concrete steps of the museum. Although the species has been reported from a Pennsylvania greenhouse (Dundee, 1974), this is the first record from a metropolitan out-of-doors site. Additional North American localities include Illinois, Nova Scotia greenhouses, and Tennessee (Dundee, 1974), San Francisco, California and Oregon (Hanna, 1966).

Native to Western Europe, Algiers, Madeira and the Canary Islands (Tryon, 1885), this is a very secretive burrowing form which spends most of its time underground feeding principally upon earthworms (Hanna, 1966).

Two additional species of *Testacella* have been reported from North America, *T. scutella* Sowerby (considered as conspecific with *T. haliotideia* by some authors) and *T. maugei* Férussac.

LITERATURE CITED

DUNDEE, D. S. (1974) Catalog of introduced molluscs of eastern North America (north of Mexico). -- *Sterkiana* 55: 1-37

HANNA, G. D. (1966) Introduced mollusks of western North America. — *Occ. Papers Calif. Acad. Sci.* 48: 1-108

TRYON, G. W. (1885) *Manual of Conchology*, 1: 1-364, 60 pls.

Accepted for publication March 3, 1976

REPRINTS OF RARE PAPERS ON MOLLUSCA: MÜLLER, 1774, VERMIUM TERRESTRIUM ET FLUVIATILIMUM, SEU ANIMALIUM INFUSORIORUM, HELMINTHICORUM, ET TESTACEORUM, NON MARINORUM SUCCINCTA HISTORIA HAVNIAE ET LIPSIAE, APUD HEINECK ET FABER

FOREWORD

Anyone interested in land and freshwater Mollusca has encountered and perhaps deciphered the cryptic 'Müll., Verm. Terr.' and soon finds out that this refers to the great work by Otto Friedrich Müller on the non-marine Mollusca of Denmark. Müller mentions the snails and clams of Europe and even gives their vernacular names but at first sight it seems incongruous to find his name attached to North American species. The fact is that his Danish species were not all confined to Europe but were holarctic in distribution and his work was useful not only to European workers but also to that small band of American malacologists who were collecting and describing American mollusks in the early nineteenth century. The need to consult Müller from time to

time still makes itself felt but it is a lucky malacologist who has easy access to Müller's book. It seemed a good idea, therefore, to reprint it serially in *Sterkiana* and to print some extra copies to bind together later on. The book has some 230 pages and is well within the space limitations of five or six numbers of *Sterkiana*.

The University of Michigan, with customary generosity, permitted xeroxing of their copy and reproduction in *Sterkiana*. This gracious gesture will place in their debt all but a few malacologists, i.e. those who own or have access to an original copy.

A. L.

VERMIVM
TERRESTRIVM ET FLUVIATILIVM,

SEU

ANIMALIVM INFUSORIVM,
HELMINTHICORVM, ET TESTACEORVM,

NON MARINORVM

SUCCINCTA HISTORIA,

AUCTORE

OTHONE FRIDERICO MÜLLER,

REGI DANICÆ A CONSILII JUSTITIÆ ET CANCELLARIÆ, ACAD. SCIENT.

NAT. CURIOS. HOLMENS. ET BOICÆ, NIDROSIENS. BEROLINENS.

ALIARVMQUE SOCIET. LITTER. SODALI. ACAD. PARIS. CORRESP.

VOLUMEN ALTERVM.

HAVNIÆ ET LIPSIÆ,
APUD HEINECK ET FABER,
EX OFFICINA MÖLLERIANA.

1774.

Qual Vitruvio fabrica loro una casa sì capricciosa, e impossibile a imitarsi dall' arte? Iddio, compreso sotto il vocabolo di Natura, geometrizza in ogni suo lavoro, come dicean gli antichi, onde possano con uqual fatica e diletto nella semplici voluta d'una Chiacciola raggirarsi i pensieri. BUON. RIGREAT.



Satis lapillis & testis lufum est; fatis structura cochlearum & concharum multiplici, earundemque coloribus in infinitum variantibus stupuimus. Nuces jam demum pueris relinquere convenit; viros nucleum, non putamina, quærere, *hospitem*, non domicilium unice, mirari penitusque noscere decet. Hunc si noverimus, rationem diversæ habitaculi structuræ, & ipsorum forsitan colorum in eo inveniremus, hallucinationesque auctorum in distributione generum evitarem.

Partes sexuales vegetabilium, in pluribus minimas, in multis dubias pro fundamento systematis admittunt Botanici, totum animal in cognitione & distributione *Testaceorum* negligunt Zoologi, testarum figuris, dentibus & aperturis, characteri in pluribus generali, lubrico, vago, unice inhærentes. Hinc *Testacea terrestria, fluviatilia, marina, operculata, & operculo destituta, quadricornia & bicornia*, quan-

tumvis maxime discordia ad unum idemque genus vel promiscue rediguntur, vel in genera Naturæ absona confuse dispescuntur. Hujus exempla apud auctores, qui animalia ignoravere, illius in HELICE perillustris a LINNÉ, exstant. Jure quidem miratus est *Fabius Columna*, neminem omnium, qui de testaceis verba fecere, in testis inhabitantia animalia considerasse, variasque eorum effigies tradidisse, at mirabundus ipse in trium descriptione subfritit. Inde centum & quinquaginta & quod excurrit, anni effluxere, antequam ullus, (anatomicas perpaucorum descriptiones non moror) nutui naturæ & monito *Columna* morem gesserit. Primus post tantum intervallum clariss. *Adanson* summa opera & sagacitate testacea in tractu Senegalensi Africæ obvia investigavit, primus celeberr. *Geofroi* in domestica inquisivit, eo animo uterque, ut ad genera effingenda structuram animalis in consilium vocaret.

Testaceis *terrestribus & fluviatilibus* agri *Fridrichsdalensis* per aliquot annos intentus, quædam nova, plura scitu digna inveni, quæ cum cultoribus rei naturalis communicare mens est. Ut vero minus cognita noscantur, notaque facilius quarantur, methodum fingere, vel jam fictam sequi necesse est. Habeant suas divisiones, qui conchylia ad pompam colligunt,

ligunt; arbitrio cujuscunque auctoris innixas; naturæ studiosi naturam consulant, naturamque sequantur, saltem in iis, quæ ante pedes sunt, & quantum licet. Methodum enim, qui ex mera testarum inspectione, nullo habito inhabitantis respectu, concinnat, *Cytherem* & *Lyncem* Monoculorum, vera insecta, in numerum testaceorum admittere, *aquatica* & *terrestria* eidem generi adscribere, hospitesque simillimos ob diversam habitaculi faciem separare debet. Qui ex structura testarum & vermium simul dispositionem componit, ex unico *Helicis* genere septem saltem genera, ex *Planorbe* tria ex *Nerita* tria, invita quidem natura, fingat. Sic clarif. *Geofroi*, qui in definitione generica testas quoque admittit, eodem jure, quo *Ancylum* a *Planorbe*, cum sola testa differre contendit, sejungit, propriumque genus constituit, *turritas* *Helices* a *depressis* & *globosis*, *depressos* *Planorbes* ab *ovatis* disjungere debuisset; at recte sejunxit, novumque genus finxit, licet veram & validam novi generis rationem minus observaverit; limax enim *Ancyli* non tantum testæ structura a *Planorbe* diversus est, sed tentaculis quoque, quæ in hoc *longa* & *setacea*, in illo *brevia* & *truncata*, sunt. Ea quidem divisio, quæ & ad testas & ad vermes respicit, adeoque & sensibus & rationi satisfacit, optima esset, si figura testæ Structuræ hospitis externæ responderet, cum vero limaces
genere

genere similes domunculas genere diversas habitent, nulla omnino ex utroque nota differentialis generica peti potest. Ordines hinc ex structura vermis generaliori; genera ex numero figuraque tentaculorum, ex situ oculorum: subdivisiones & species ex figuracione unicuique testæ propria depromsi. Hoc Naturæ maxime congruum esse, ex eo patet, quod animalia simillima, externaque facie prorsus eadem, vel nulla vel testis infinite variantibus, lege tamen omni generationi immutabili, instruantur, dehinc ratio, ob quam limax *nudus* & testa *tectus* eidem ordini, *terrestris* vero, *fluvialis* & *marinus*, testæ licet simillimæ sint, diverso generi submitti debent, ipsis oculis percipitur.

Objici quidem potest, testarum hospites explorari in folis domesticis concedi, at ex eorum inquisitione tanto plus voluptatis & emolumenti redundaturum, quanto opus quodlibet artificiosissimum opitice inferius est, abunde evincunt observationes in oeconomia horum animalium nuper institutæ, exoticorumque limacum investigationem, crescente indies in Historiam naturalem favore, Conchyliologis peregrinantibus curæ cordique futuram, ominantur. Hac ratione, si unquam, Systema Conchyliorum naturale sperari potest. Difficillimum quidem argumento
geminò,

gemino, at minus justo arguit clarif. *Schröter*, quod nempe *testacea Gallia in Germania non æque indigenæ sint*, quodque *differentia hospitiæ testarum generis diversi nulla, vel saltem indistincta sit*; hoc maxime probat, testas similiarum animalium, licet sint diversæ figuræ, minus tamen bene diversi generis dici, cum a potiori sumi debeat character genericus, ac similia animalia, ædificia etiam si diversa inhabitantia, uno nomine jungi, dissimilia, inhabitationes quamvis similes sint, separari e nutu Naturæ; illius vero contrarium verum est, testacea enim fluviatilia & terrestria nulli regioni propria, sed maximam partem totæ Europæ communia sunt.

Quibus interim testarum contemplatio sufficit, divisiones a figura domicilii ubique apud auctores reperiunt. Ego, loco taliscunquæ distributionis arbitrariæ, contra primam operis ideam, quæ circa indigenarum historiam unice versabatur, utque historiam ostragodermatum *terrestrium & fluviatilium* *) quam maxime completam traderem, quascunquæ testas, quæ vel aucto-

ritas

*) Divisio Testaceorum in *terrestria & fluviatilia*, si testas respicis, naturæ repugnat, si vero animalia, maxime consentanea est; terrestribus enim non tantum tentacula forma & structura a fluviatilibus diversa, sed iis quoque solis quatuor tribuit, nec paucissimorum utriusque classis in aquis & in terra momen-

ritas descriptorum, vel habitus terrestres & fluviatiles fecit, in Museo proprio vel civium obvias descripsi, suoque loco inserui. Hoc quidem ex parte & ante me præstitere viri illustres; *Geofroi* enim terrestria & fluviatilia, *Schröter* sola terrestria, *domestica* descripsere, *Lister* vero & *Martini* de tota Testaceologia optime meriti, quicquid de utroque cujuslibet regionis eousque innotuerit, ex propriis aliorumque observationibus collegerunt. Qui tamen in hoc opere numerum novarum & minus notarum specierum computaverit, descriptiones omnes de novo factas, observationes quam plurimas, erroresque auctorum paucis castigatos viderit, nihilque aliorum auctoritate, sed quodvis, minus vulgare, propriis observationibus prolatum, observationesque auctorum meis non contrarias, ne centies dicta regerantur, omittas animadverterit, operam supervacaneam minime censeat, viris præclaris publice & privatim editionem operis exoptantibus gratias habiturus.

Specialiora de domunculis & inquilinis testaceorum in ipsa cujuslibet historia monui, generalioribus hic locum servavi, cumque intellectus humanus angustis finibus

terrenam hospitiū terrenam gentem in aquaticam, vel hanc in illam mutat; ægrius *marina* a *fluviatilibus* separantur, at hoc haud præsens indaginis est.

finibus circumscriptus sit, ac memoria facile labet, hallucinationes sagacissimis quoque non possunt non obrepere, eo vero exactiori cura dispellendæ sunt, quæ majori auctoritate derivantur; paucis igitur, quæ circa testacea terrestria & fluviatilia versantur, nec in ipso opere memoratæ sunt, notare lubet.

Limacès quadricornes omnes terrestres sunt, minus vero omnes terrestres quadricornes; *Vertigo* & *Carychium* legem hanc Conchyliologis generalem limitant. Aquatici omnes *bicornes*, minus vero vice versa. *Helicem succineam* auctores *amphibium* dixerunt, quum ei soli proprium in aqua æque ac in terra vivere crederetur, at hoc pluribus *terrestrium* commune est, multos enim aquæ immixtos non suffocari, quosdam sese, ut aufugiant, aquæ sponte tradere, ideoque locum aquis clausum, quem cochleariis instituendis *Varro* indicat, non satis tutum, varietatemque *H. nemoralis*, quod singularissimum puto, fundo rivi tota æstate vivere, observationibus didici. *Fluviatiles* non contra: aqua enim extracti & in sicco positi brevi pereunt; notandum enim *H. succineam*, quam fluviatilem censent plerique, esse terrestrem, id quod tentaculis quatuor & saliva, qua aperturam more terrestrium claudit, egregie probatur. Plures tamen aquam sponte deserunt, plantis aquaticis vel parietibus

vasis, quo continentur; extra aquam sese affigunt, ibique post aliquot horas plerumque pereunt.

Terrestrium & fluviatilium *victus* ex herbis est; nudi Fungos, Boletos & Agaricos inprimis in deliciis habent; at inediam ultra fidem pati possunt; miratur quidem *Lesser* decemdialem, ego vero Hel. *nemorales* ultra annum absque omni nutrimento, ac *Buccina* & *Planorbis* per integros menses in aqua immutata omnisque vegetabilis experte, vivos servavi; fluviatiles guttulas aquæ, hisque immixtas moleculas terrenas, animalculaque microscopica fere continuo ingurgitare vidi, terrestres vero solo humore aeris, pluviaque naturali vel factitia, quam, uti plantæ marinæ aquam, poris corporis undequaque hausisse suspicamur, vitam sustentare. Errant tamen, qui testacea sola humiditate aquarum ubique in poros penetrante nutriri putant; organa oris nec terrestribus nec aquaticis frustra tributa sunt, utraque nutrimentum ore capere sæpius vidi. Papyro quoque & calce vesci dicuntur, mihi vero *Limaces* & *Helices*, ut sese carcere liberarent, chartam rodebant & dilacerabant, *lapicidamque* auctorum ab omni calce remotam vitam degere infra dixi; quo medio panem, caseum & piscium caudam salsamque

*) Hinc effatum Listeri & aliorum *cocleas terrestres marino & dulci aqua aque necari*, anatom. p. 81., in genere non valet.

que carnem e longinquo olfacient, unde ipse *Lister* hoc sciverit, ignoro. Fluviatilia ore & testa hiante guttulam roris appetere, sacula majorum sapit.

In terrestribus decantata poetarum & pictorum, quibus quælibet audendi fingendique potestas, *sagitta* amatoria revera existit; primo vere ubique in hortis inventu facilis, perpauca licet mortalium unquam visa sit. Conchyliologorum princeps *Lister* eam jam vidit, obiterque delineavit; corpore *H. nemoralis* exitus infixam sapius extraxi, amicosque extrahere feci. Candida est, pellucens, cartilaginea, quadrata angulis submembranaceis, apice acutissima, basin versus aliquantum coarctata & teres, ipsa basis vero orbiculata est, & truncata: latera percurrit quasi linea utrinque pinnata; lineam lata, octo lineas longa est. Curvam minus bene *Lesser* dixit, cuspide quadruplici instructam perperam anonymus *). Humoris cærulei, quem ex vulnere defluere narrat *Lister*, nec vulneris ullum vidi vestigium, licet ex utroque in ipsa copula sagittas extraxerim.

In testa tectis coitus non ante peragitur, quam testæ incrementum absolutum sit; absolvitur enim in plerisque, cum limax marginem aperturæ labro firmat. Hoc pubertatis indicium, jam enim materies, quam natura exstruendæ domunculæ suppeditat, formationi sa-

b 3

gittæ

*) Schauplatz der Natur, vol. I. p. 277.

gittæ cupidinis, propagationi & organis ad eam necessariis impenditur. Quoties coire cupiunt, altera alteri sagittam vibrat, hæc vel juxta decedit, vel alteram tangens corpori fixa inhaeret. *Helices nemorales* versus finem Aprilis & Septembris, medio Junii & Augusti, & sub initium Maji copula junctas deprehendi.

Varietates *Limacis cinerei* ad meridiem & die serena ineunte Augusto *Lister* in copula vidit, ego eodem anni tempore, post meridiem quoque & tempestate maxime pluvia & ventosa individua varietatis *Limacis atræ* & individua variet. *L. cinerei* mutuo junctæ contemplatus sum. Ope humoris viscerati & indurati, caudas involventis, corpora utriusque, triplici gyratione implexa, tres ulnas supra terram de trunco fagino libere pendebant; infra capita a dextra colli parte propendebant genitalia utriusque glutinosa septem gyris, plicis innumeris, circumacta, motu serpentiformi a dextra in sinistram et v. v. continuo alternante; oræ alter alterius foramini anali adjacens, glutinosam genitalium substantiam mordens suctu alternatim adtraxit & evomit, suavisque ejusmodi speciem propagabat. Foramen *genitale* ab *anali* diversum est; hoc in ipso clypeo limacis nudi, in collari testa recti, respirationi simul inserviens, illud pone caput, utriusque in dextro latere; quod monendum erat, quia illustribus a *Linne* & cl. *Bomare* utrumque, quod Natura satis
removet;

removet, auctoresque apprimè distinxerunt, unum idemque dicunt; *Argenville* *) vero anum & exitum excrementorum diversis locis oscitanter posuit.

Limaces sub lapidibus & intra cortices arborum emortuarum &c. sese Octobri condunt, hisque locis duos & saepius plures haud tamen diversæ speciei in societate constipatos tota hieme latuisse reperi. Caput clypeo condi potest, minus vero cauda & totum corpus.

Tentacula objecto tentando vix inserviunt, nec iter dirigendo, uti volunt auctores **), limaces enim tentaculis orbati æque libere & celeriter repunt; at sensu nobis ignoto forte gaudent. In *quadricornibus* ope nervi nigri, quem opticum vocant, in sese revolvi possunt, & quidem in nonnullis maxima celeritate ad medium dorsi usque; in *bicornibus* seu aquaticis ope musculorum vel rigida extenduntur, apiceque recurvantur, vel laxa dependent, vel ad corpus reflectuntur, more vero terrestrium contrahi & in capite condi, quod

*) Zoomorph. p. 87. leur anus s'ouvre à la partie droite du colier & leurs excréments sortent par un trou voisin.

**) *Argenville* Zoomorph. p. 87. idem oculos, quos limacibus testis tribuit, nudis adimit, ibid. p. 84 & olim *Plinius*: *cornua prozendentis, contrahentesque oculis carens, ideo corniculis præsertim iter*; h. nat. l. 9, c. 32.

quod indicat LISTER, minime possunt; *oculi* aquaticorum vestiuntur membranula pellucida, quam limax, ut objecta perspicue lustret, in colliculum erigit. Tentacula *inutilia & superflua* temere pronunciat clarif. ADANSON, cum ab ignorantia usus ad non usum minus logice concludatur; si limaces tentaculis orbatos vitam æque degere, ac iisdem instructos cognitum habuisset, specie probabilitatis sese commendaret nimis jam audax assertio, quid vero, si toto capite orbi vitam ultra annum producant, an caput igitur inutile & superfluum? cura, qua Natura capitis & tentaculorum restitutionem molitur, sensusque exquisitior quam in reliquo corpore, quod ipse eodem loco confitetur, si quodvis aliud argumentum deesset, hæc organa æque utilia & necessaria esse, evincit.

Membrana, quam veteres *limbum*, hodierni in limacibus seu cochlearum inquilinis *collare* sive *pallium* nominant, collum animalis cingit, ac ex sententia quorundam testam interne & externe in nonnullis vestit. Usus ejus secundum auctores varius: hiatus duos, alterum foecibus, alterum aeri inservientem continet; ADANSON putat, pallium, collare, ne aqua animaculo invito testam intret, impedire, vel intromissam retinere. An limax unquam aquam in interiora testæ admittat, incertus sum, nec video, collari aper-
ruram

turam occupante, quomodo fiat, nisi foramine vel branchiali vel annali, hoc vero, dum foeces excernuntur, modo aperitur, illud satis patulum spatium vastum seu interiora ipsius testæ ab omni aqua vacuum parietesque interiores membranula glutinosa obtectos contemplari permittit. LISTER quidem meminit, *Buccinum foramen ad dextrum rotundum & angustum sibi efficere solere, quo piscium more quodam reciproco aquam recipiat ac ejiciat.* Foramen dextrum millies inspexi, aquam intrare & ejici in *Buccino* nunquam vidi; *Planorbis* quidem & *Neritæ* e laxo pallio siphonem seu tubulum, huic forte usui accommodatum, quandoquidem formant, nec tamen in his usum aquæ reciprocum percepi. Membranam triplicem glutinosam distingo, mediam & crassiolem, duasque graciliores, mediæ utrinque adnatas. Illam *Collare*, has *Tunicam* & *Pallium* dico.

Collare est membrana opaca, & spumosa, qua limax cingitur & pro lubitu conditur, aperturam testæ, si vel limax domunculo exit, vel in ea latet, continuo occupans & implens. In eo, limace foris egresso, duo foramina observantur, trachæ & ani; hujus ope excrementa secernuntur, illius aër spiratur; utrumque pro lubitu animalis aperitur & clauditur. Præcipuus ejus usus & nunquam non mirandus, in ter-

restribus spumam, ex qua opercula momento formantur, secernendi, in omnibus testam seu domunculam struendi, laesam refarciendi. *Tunica* est pellicula, venis, maculis, punctisque variegata, parietes interiores anfractuum obvestiens, ac colores, quibus testæ animalculo foetæ superbiunt, intrinsecus suppeditans. Huic glabrities interior testæ, huic (minime corpori limacis, quamvis maculatum sit) omnes testarum maculæ & puncta aurea, nigra, lutescentia &c. ac venarum ramificationes per testam apparentes, in plurimis nostratium terrestrium & fluviatilium debentur; licet enim foris egrediatur animal, maculæ & puncta in testa permanent, ac si domo clauderetur; e contra, cum sese in anfractus minores subducit, maculæ & puncta in majori evanescunt, & si totus extrahatur, tota testa immaculata evadit, sola fasciæ in fasciatis testæ connata superstite. Per foramen branchiale *collaris*, quod aperitur, quoties limax foris sit, interiora anfractus majoris lustrare, lateraque undique *Tunica* reticulari cooperta contemplari licet. *Collari* & *tunica* omnes cochleæ instruuntur, *Pallio* vero perpaucæ fluviatilium tantummodo *) gaudent. Hoc vel integrum, vel in lacinias fissum superficiem testæ vel

*) In terrestribus mihi obviis sola *Helix pellucida* p. 16. rudimentum pallii exserit, laciniam nempe, qua testam perpolit. Ibidem & in schemate generali collare ex sententia auctorum pallium dixi.

vel ex parte vel totam, quoties limax foris vage-
tur, obtegit.

Licet limaces toto corpore *humorem* exsudent, tria tamen loca, quibus præcipue fecernitur, notari merentur; *collare* nempe, de quo jam diximus, pluraque infra dicentur, *fossula* triangularis in dorso juxta extremitatem caudæ, e qua viscus momento coagulatus exstillat, novusque, quoties acicula auferatur, brevi succedit, & ipsa corporis *extremitas*, ubi alius liquor viscidus, quo limax vestigia sua notat, excernitur. In tanta humorum affluentia Limacis terrestris ingressus a LISTER apprime dicitur *quædam veluti natiatio in proprio humore*, perverse vero subjicitur: *si non continuo illum inter eundem eructaret, propria saliva implicitus loco adfigeretur*, dicere potius voluit, si inter eructandum cursum non pergeret, nec hoc satis justum, limaces enim facultate hunc humorem pro lubitu fecernendi vel retinendi gaudent, alias enim, cum sæpe extra testam temporis spatio quiescant, ibidem rore pingui coagulato tenerentur. Spuma, quam lacestræ cochleæ copiose eructant, non e foramine branchiali, uti Lister narrat, sed ex poris *Collaris* enascitur, ex ore etiam more *Larvarum* salivam, cui minus bene formatio fimbriæ testarum tribuitur *), evomere, at rarius, vidi.

c 2

In-

*) Klein test. form. p. 16 & 24.

Incessus limacis ope organorum in area media plani inferioris corporis latentium peragitur; ibi enim in nudis præsertim motus undulans concitatus, continuo sursum tendens, id est, ab extremitate pedis postica ad anticam, clare conspicitur. Lente tamen loco in locum movetur, etiamsi undarum altera alteram celeri lapsu cogat. In fluviatilibus nulla quidem undulatio percipitur, ope tamen occultæ rotationis vel ignoto mechanismo nec lentius, quam terrestres, progrediuntur.

Limaces terrestres testa instructos imprimis destruunt *Lacertæ* *) & *Staphylini*, fluviatiles vero *Hirudines*, carnem limacinam exfugere appetentes.

Ætas & numerus annorum nec ex numero anfractuum nec ex productione aperturæ in cochleis judicatur; ex illo quidem juniores & ætate proveciores distinguuntur, nulla vero annorum determinata mensura sumi potest; incrementum enim testæ, quod fit novi succi indurati appositione annua ad marginem aperturæ, ratione ætatis, tempestatis, valetudinis, nutrimenti &c. variat, cochleaque ad justam magnitudinem producta, vel potius limace generationi maturo,

*) Jam Plinii tempore lacertæ inimicissimum genus cochleis. *H. nar.*
l. 8. c. 39.

turo, testæ margo in genere non amplius increfcit, sed in terrestribus saltem labro terminatur. Falsum dehinc quorundam judicium, cochlides totidem annos, quot orbis habere; falsum & contradictorium terrestres numero anfractuum octenario constanter gaudere, ætateque ad decennarium augeri. Numerus in perfectis quinaris vulgaris, binarius, ternarius & sedecennarius rarus, intermedii passim obvii; quilibet suæ speciei plerumque constans.

Rationem plurimarum testæ circumvolutionum in eo minus bene *Lister* quærit, quod limaces testis non sint affixi, corpusque eorum dehinc, ne excidat, multiplici gyro intorquatur, ligamentis enim parieti testæ interiori adeo adhærent, ut cadavera eorum non nisi violenta ligamentorum ruptione educantur. Quomodo *Klein* duas priores spiras cochleæ junioris minores duabus prioribus cochleæ adultæ dicere queat, non video, in omni enim ætate volumine eadem & immutabiles persistunt, nec ullo modo augeri possunt vicinis anfractibus stipatæ; hanc ob causam, quoties de specie juniorum seu imperfectarum cochlearum ambigitur, diffractio seu reductio perfectæ, cujus pullos suspicaris, ad totidem anfractus, omne dubium tollere valet.

Apertura cochlearum in quibusdam semper patet, in aliis *operculo* clauditur; in *Buccino* & *Planorbis*

continuo, in *Helice*, *Vertigine* & *Carychio* pro tempore pater, interdum operculo *factitio* obturatur, in *Nerita* & *Valvata nativo* clauditur. Hoc dehinc perenne pedi limacis adhæret, ac pro ratione aperturæ novis accessionibus augetur; illud a corpore disjunctum improprie operculum dicitur, pro lubitu animalis, effingitur & destruitur, æstivo tempore membrana pellucida simplici constat, momenta vel dies perdurat; autumnali vero membrana triplicata, calcarea, opaca, totam hiemem duratura, componitur.

Operculo *factitio* aperturam pro lubitu claudendi facultate omnes *Helices* mihi notæ gaudent, *Turbini*bus cl. *Schröter* nequidem exceptis, in *Hel.* enim *lubrica*, *obscura*, *peruersa*, *Trochulo* &c. operculum tenue & pellucidum, at non calcareum, paucissimis proprium, vidi.

Colores testarum alii epidermidi, alii ipsius testæ substantiæ calcareæ deberi dicunt; in genere huic debentur, illamque omni pictura carere, fasciasque & maculas diversicolores ipsi substantiæ testarum coalitas per epidermidem pellucere, in quibusdam terrestribus & fluviatilibus reperi. Aliam colorum rationem infra exemplis demonstrandam, supra indicavi: *Tunnicam* nempe limacis testas pellucidas & immacu-

tas

tas punctis, maculis, venis quoque ramosis varie & graphice exornantem. Enumero & diverso fasciarum situ characterem specificum perperam desumi in ipsa testaceorum historia luculenter patet.

Cum de distributione & partibus conchyliorum multa differuere auctores, paucula de binis maximi ponderis momenti in historia testaceorum inter gravissimos Naturæ scrutatores huc usque controversis modo interponere libet. De *formatione & incremento* testæ, ac de *restitutione* partium amissarum in limace res agitur.

Quoad primum duplex est sententia Physicorum, altera nempe *animalia testacea cum ipsis & per omnes dimersiones integris testis ex ovo in lucem prodire*, altera *hospites quidem ex ovo at non una cum testis numero anfractuum perfectis enasci*.

In spermate testaceorum, imprimis fluvialium & marinorum, sæpenumero vidi, nudis quoque oculis facillime conspicitur, limaces ibidem jam testis reptos latere, testamque & hospitem, dum genituræ pellicidæ adhuc claudantur, æque ac extra eandem libero aëre, incrementum sumere. Hoc facilis anfractuum,
in

in junioribus & adultis diversa *), computatio probat; testa enim cum spermate excludatur, in *cochleis* uno vel binis anfractibus instructa, in *conchis* paucillo stratum numero circumscripta est, adulta vero ætate in his strata fere innumera, in illis quinque, octo & plures circumvolutiones observantur. Hinc dubitari vix potest, testam, ut primum limax evolvi coepit, ratione hujus voluminis simul incrementum, utrumque successu temporis ad magnitudinem speciei cuique determinatam augeri.

Modus, quo incrementum in testa perficitur, æque ambiguus habetur. Res augetur vel introrsum evolutione in majus volumen, vel extrorsum novis additamentis; illud augmentum *per intussusceptionem*, hoc *per juxtapositionem* nominaverunt; hanc in testis formandis *Reaumur* adoptavit, observationibusque probavit; *Klein* vero sententiam Reaumurianam longa dissertatione impugnans, intussusceptionem in testis æque, ac in hospite admisit. Quilibet affectus suos habet, lisque ad hodierna usque tempora sub iudice persistit, quam cum novis observationibus dirimi posse

*) Contrarium quidem disputat *Klein*, cochleam nempe in ovo tot spiras habere, quot habeat ad profectam ætatem devoluta, at illusionem hanc facile dissipat quælibet junioris & adultæ testæ ubivis obviæ inspectio.

posse judicarem, non absque miratione audio, illustrem BONNET experimentis claris *Herissant* circa ossium & testarum generationem persuasum, se ipsum amicumque suum REAUMUR erroris accusare *) ac in sententiam *Kleinianam* descendere. Tanta tamen auctoritate me minimum convictum profiteor; claris enim *Herissant* experimentis summa industria institutis, testas conchyliorum duabus substantiis, membranacea seu animali & terrea seu cretacea constare, hancque illam incrustare, tantum probat; addit quidem, testam esse appendicem corporis limacini, fibrisque ligamenti, quo limax testæ adhæret, continuari, hypothésin tamen nec argumentis aut observationibus firmat, nec *Reaumuriana* contraria refutare conatur.

Non parum lucis ex operculis Helicum factitiis, examini nitri *dissolventis* nominati auctoris subjectis, sperabam; collare enim Helicum exsudare liquorem, in operculum indurescens, hocque semper liberum & absque ulla cum limace connexionem aperturam testæ claudere

*) *J'avois donc commis une erreur sur les coquillages, & cette erreur je l'avois commise d'après feu mon illustre ami Mr. de REAUMUR; des expériences équivoques l'avoient trompé; la coquille ne croit point par apposition ou par transudation. Paling. philos. t. 1. p. 405.*

claudere sæpius vidi. Concludebam hinc, si ope nitri *dissolventis* meram substantiam cretaceam, nullam membranaceam, reticularem; qualis in ipsa testa adest, deprehenderem, clarif. virorum *Klein & Herissant* hypothesin de incremento testarum per intususceptionem veritati proximam esse, si vero membranacea, seu animalis, in operculo obvia esset, hanc æque in testa ac in operculo vaporum *collaris* exhalantis debitam referri, adeoque incrementum fieri per appositionem. Hæc meditati occurrit ipse desideratus Galli commentarius *); non minore voluptate membranam animale operculi *H. pomatie* nitidissime impressam lustro, cumque nullam operculi cum limace cohesionem unquam viderim, aut suspicari licuerit, sententiam de incremento per appositionem non parum corroboratam censeo; averfari quidem videtur ligamentum vasculosum, quo operculum pedi limacis adhærere afferit auctor, ingenue tamen, licet fidem dictis clarif. Viri detrahare nollem, confiteor, me nullum ligamenti vestigium unquam in pede limacino, liquorem vero undique e *collari* spumantem paucis momentis in operculum evadere sæpius vidisse, nec quo modo ope unius ligamenti seu canalis advectitii, & quidem non centro operculi sed lateri affixi operculum adeo compositum, quale

*) Memoires de l'Academie des Sciences; à Paris 1766.

le in *H. pomatia*, brevissimo temporis intervallo construatur, conciliari posse. Valde quoque dubito, observatores, si illustrem *Reaumur* excipias, operationi limacis in extruenda testa, aut operculo, interfuisse, alias enim nec corpus limacis archetypum testæ seu gyrationum, nec e liquore viscido corporis testam, nec spumâ oris operculum & labium formari dixerint, cum formatio hæc in Helicibus, incrementum & refarcitio testæ in omnibus, soli *collari* deberi videtur. Spumantem e collari liquorem in membranam operculi diaphanam mutari in terrestribus cochleis, inque *H. pomatia* & aliis incrustari, cuilibet videre est. Ipse *Klein* docet, maximum cochlearum anfractum appositione augeri, limbumque seu labium saliva oris (*vel potius* collaris) eructata redintegrari, cur non æque minores tali additione increfcere possunt?

Testarum & operculorum partes constituentes, substantias nempe membranaceas & cretaceas optime extricavit clarif. *Herissant*, at utriusque generatio ope juxtapositionis aque ac intussusceptionis peragi posse videtur; illa in formatione operculi ad oculum demonstratur, hæc sola cohæsiõne ligamenti limacis cum testa inicitur; hoc tamen, cujus unius medio incrementum per intussusceptionem, si unquam, fieri potest, nullam canaliculorum limacis & testæ liquores nutritios vehen-

tium communionem admittit, quoniam locum ad perfectionem testæ usque mutans, parieti novo & politissimo sensim affigitur. Hæc ligamenti translocatio absque controversia est, quicquid dicat *Klein*; in pluribus enim cochleis pellucidis anfractus minores a limace derelicti vacui videntur, *Helices*que apice *Planorbis*que centro mutilatos vivos servavi; hinc nec, uti asserit *KLEIN*, *ab illo puncto primæ spiræ ejusque salute vita animalis ejusque vegetatio dependet*; nec causa, cur prima & secunda spira pertusa non refarciatur, in tenerrimis animalis interaneis quærenda, sed in remoto collari, reparationem unice efficiente.

Quoties limax domunculam nova accessione ampliare molitur, testam non operculo claudit, nec sese in interiora subdit, sed collari involutus in ipsa apertura dies immotus hæret; tum in parte collaris, quæ oræ aperturæ affixa aliquantum prominet, porisque referta est, humor nitidus, ac ope lenticulæ undulatio molecularum percipitur; ejusmodi succi testificantis exhalatione lamella altera alteri substituitur, incrustatur, testaque augetur; pars novitia veteri mollior, & flexilis aliquamdiu persistit: reparationem fracturæ e contra tentans fragmента primum removet, dein collari parti læsæ interiori affixo, lacunam membrana opercula claudit, ita tamen, ut ora collaris, restitutionem operans, libero aëri exposita sit. Sententia dehinc de incremento

cremento testarum per juxtapositionem, immortalis REAUMUR meisque observationibus nondum refutatis, facta recta persistit, phaenomenaque allata ampliore inquisitione dignissima, naturæ consultis commendanda.

Præter hanc testarum amplificationem & restaurationem omnibus communem, limax terrestrium ætatem adultam adeptus marginem aperturæ fimbria, quam *labrum* appellamus, coronidis loco in plerisque firmat; hoc in paucis dentibus armatur, foramen axis in perforatis obturat, umbilicum vero in umbilicatis vel ex parte vel neutiquam tegit. Labro ita ulterius testæ augmentum cessat; in paucissimis tamen, rationem ignoro, novum incrementum insequitur, secunda & tertia *) coronis imponitur. Mutationum harum inscitia Conchyliologos in errores abduxit, cochlearum enim juniores & adultas, perforatas & imperforatas, labiatis & labri expertes, dentatas & muticas specie diversas descripserunt, cum cujuslibet animantis & vegetantis character specificus a perfecta & adulta ætate petendus sit, quæ cochleas muticas passim dentatas, perforatas maximam partem imperforatas sistit, umbilicatas *) vero immutatas monstrat.

d 3

Alterum

*) Hinc suturæ, unde auctores sæpe notas specificas sumere, plerumque accidentales & merum labrum relictum denotantes.

**) Cochleam *umbilicatum*, si in cavo axis anfractus conspicui sunt, sin minus, *perforatam* dicere oportet.

Alterum, quod nostra ætate historiæ naturalis cultores maxime attentos dubiosque tenet, *partium amissarum* in limace in integrum *restitutio* est. Apparitio hæc singularissima naturæ consultos memoris notæ Italos, Gallos, Germanos confestim, & jure quidem, totos occupavit, detruncatioque generalis genti limacinæ imminere videbatur; pruritus vero multorum strage inclarescendi opinione citius deserbuit, successusque diversus plurimos ambiguos reliquit. Redintegrationem affirmant clarissimi viri *Spalanzani*, *Schaffer*, *Lavoisier*, *Rose* &c. negant æque clari *Cotte*, *Bomare*, *Schröter*, *Wattel* &c. quivis pro eventu tentaminum; minori tamen jure negatur, centum enim experimenta varias ob causas frustra suscepta, unam veram & justam observationem tollere minus valent. Quam primum rumor Italici facinoris ad me venit, idem in nostratibus limacibus tentare aggredior, eventumque Academiæ Reg. Sc. Paris. anno 1769 commentario mandavi; præcipua ejus confectaria heic brevi sistere instituti ratio jubet:

Quoties caput limaci amputaverim, in detruncato capite mihi ipsi & præsentibus tentacula porrecta, puncta bina nigra, succum nervi optici, maxillamque monstravi.

Vulnus

Vulnus motu undulatorio continuo partium vicinarum intra pauca temporis minuta adeo diminuebatur, ut fere clausum videretur.

Partes integrascentes a veteri corpore candore & pelluciditate clare distinguuntur.

In individuis *Helicis Pomatie*, *Hirudinis Sanguifugæ* & *vulgaris* nulla redintegratio successit, quælibet enim intra quindecim dies periere.

In *Hel. nemorali* $\gamma\gamma$ tentacula ad medietatem abscissa intra annum nullum profus incrementum cepere, ipseque limax tanto temporis intervallo absque victu vixit.

In *H. nemorali* ϕ , cui die 9 Julii 1768 caput totum, partem colli, anticumque pedis uno ictu detruncaveram, redintegratio lente quidem, at ex voto successit. 12 Julii centrum vulneris tuberculum flavum occupaverat. 16 Septembris sola pedis antica pars restituta videbatur. 10 Octobris, quo ægrotam meam arculæ immisi, rureque mecum in urbem duxi, omnia in eodem statu erant. 23 Martii anni 1769 tota hieme derelictam & mortuam habitam in aprico fenestræ posui; testa quidem intra dimidiæ horæ spatium

tium contra omnem opinionem exiit limax decollatus, at ne lentis quidem ope ullum redintegrationis indicium inveniebatur; licet jam omnis spes restitutionis deficeret, hospitem tamen, qui ultra octo menses mecum vixerat, servandum censui, ut viderem quousque vitam absque capite & nutrimento producere valeret. 19 Maji bina tubercula, vera majorum tentaculorum rudimenta, candida & pellucida absque ulla nervi optici nigredine conspiciebantur. 1 Junii tentacula dicta in integrum restituta erant, sinistrum tamen dextro brevius fuit; nodulus apicis, punctum nigrum, nervus niger opticus, motusque retractilis & flexilis in tentaculis solitus æque conspicuus. 29 Junii novum caput, nova colli pars, anticum pedis, tentaculaque majora æqualia candore & pelluciditate primo aspectu distinguebantur, oris vero, labiorum & minorum tentaculorum nullum adhuc indicium in hac aderat. In alia vero, cui caput die 14 Septembris 1768 detruncatum fuit, labium superius 29 Maji insequentis anni formatum erat, rimaque oris solitaria in conspectum subiit.

In *Hel. nemorali* decollata 14 Septembris 1768, nullum redintegrationis indicium 30 Aprilis 1769 apparuit; die vero 19 Maji latere dextro excrescentia quædam, cujus in medio punctum nigrum conspicebatur,